

**PHG Needs Assessment Calculator****Bhutan****Down's Syndrome**

Welcome to the PHG Health Needs Assessment Calculator for Down's Syndrome. The contents of this file are listed below.

Full name of the sheet	Short name
Country demographic, maternal health and socioeconomic indicators	Demography
Country health-service data	HealthServices
DOWNS Epidemiology 1.1: Country epidemiology	DOWNS-E1.1
DOWNS Epidemiology 1.2: International comparison	DOWNS-E1.2
DOWNS Epidemiology 2.1: Data on affected pregnancies: Research studies	DOWNS-E2.1
DOWNS Epidemiology 2.2: Data on affected pregnancies: Surveillance	DOWNS-E2.2
DOWNS Epidemiology 2.3: Data on affected pregnancies: Other sources	DOWNS-E2.3
DOWNS Epidemiology 2.4: Summary of affected pregnancies	DOWNS-E2.4
DOWNS Epidemiology 2.5: Sub-population variation in affected pregnancies	DOWNS-E2.5
DOWNS Epidemiology 3.1: Mortality data: Research studies	DOWNS-E3.1
DOWNS Epidemiology 3.2: Mortality data: Vital registration data	DOWNS-E3.2
DOWNS Epidemiology 3.3: Mortality data: Other sources	DOWNS-E3.3
DOWNS Epidemiology 3.4: Summary mortality estimates	DOWNS-E3.4
DOWNS Epidemiology 3.5: Sub-population variation in mortality	DOWNS-E3.5
DOWNS Epidemiology 4.1: Population prevalence: Research studies	DOWNS-E4.1
DOWNS Epidemiology 4.2: Population prevalence: Other sources	DOWNS-E4.2
DOWNS Epidemiology 4.3: Summary of population prevalence	DOWNS-E4.3
DOWNS Epidemiology 4.4: Sub-population prevalence variation	DOWNS-E4.4
DOWNS Interventions 1: Effect of maternal age	DOWNS-Interv1
DOWNS Interventions 2: Effect of prenatal screening and diagnosis and pregnancy termination	DOWNS-Interv2
DOWNS Needs assessment 1: Quantitative baseline	DOWNS-NA1
DOWNS Needs assessment 3: Quantitative assessment of interventions	DOWNS-NA3

(There is no sheet DOWNS-NA2.)

**Bhutan****Shared Data****Demographic, maternal health and socio-economic indicators**

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Demography information from that Calculator into here. The information should be the same.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Number of persons by age-group and sex	Estimates			Your estimates			Chosen estimates		
Age group	Male	Female	Total	Male	Female	Total	Male	Female	Total
0-4 years	31489	31064	62553			0			0
5-9 years	35547	34852	70399			0			0
10-14 years	38728	38279	77007			0			0
15-19 years	37504	37732	75236			0			0
20-24 years	40254	30320	70574			0			0
25-29 years	31386	25972	57358			0			0
30-34 years	23208	19598	42806			0			0
35-39 years	21124	17605	38729			0			0
40-44 years	16022	13878	29900			0			0
45-49 years	14895	12767	27662			0			0
50-54 years	11779	10268	22047			0			0
55-59 years	8764	7628	16392			0			0
60-64 years	7564	7010	14574			0			0
65+ years	15331	14414	29745			0			0
Total	0	0	634982	0	0	0	0	0	0
Female population aged 15-44 years		0			-			-	
Data year	2005 reported in 2005								
Source, Year	UN 2011								

**Ethnicity.** Please enter data for the main ethnic groups if you are working with a population that is different from that of the country.

Ethnic group	Number	% population

<b>Fertility and mortality</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Crude birth rate: live births (LB) / year / 1000 population	19	Unicef, 2007				
Still birth rate (SB) per year per 1000 total births	22	WHO, 2009				
Total births in 1000s (LB+SB) per year	12	Unicef, 2007				
Infant mortality rate: infant deaths / 1000 LB / year	44	UNICEF				
Under-5 mortality rate: U5 deaths / 1000 LB / year	56	(2011), 2010				
Percentage births in women >35 years						
Life expectancy at birth (yrs)	63	WHO, 2009				
% of marriages consanguineous						

<b>Maternal health</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Prenatal visits – at least 1 visit (%)	88	WHO, 2007				
Prenatal visits – at least 4 visits (%)						
Births attended by skilled health personnel (%)	71.5	WHO, 2007				
Contraception prevalence rate (%)						
Unmet need for family planning (%)						
Total fertility rate	2.6	WHO, 2009				
% home births						
% births at health care services						
<b>Newborn health</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Number of neonatal examinations by SBA / trained staff						
% neonatal examinations by SBA/ trained staff						

<b>Socio-economic indicators</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Gross national income per capita (PPP int. \$)	4820	WHO, 2008				
% population living on < US\$1 per day	26.2	WHO, 2003				
Birth registration coverage (%)						
Death registration coverage (%)						

LB = live births

PPP = purchasing power parity

SBA = skilled birth attendant

**Bhutan****Shared Data****Health Services Data**

**Please read first!** If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Health Services information from that Calculator into here. The information should be the same.

This section provides health-service-related information for your country.

**By default, the Toolkit contains information at the national level.**

**If you would like to use a different population, then replace country information with that of your specific population of interest.**

<b>Health Expenditure</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Per capita total expenditure on health (PPP int. \$)	274	WHO, 2009				
Total expenditure on health as percentage of GDP	5.5	WHO, 2009				
Per capita government expenditure on health (PPP int. \$)	224	WHO, 2009				
External resources for health as percentage of total expenditure on health	7.6	WHO, 2009				
General government expenditure on health as percentage of total expenditure on health	81.9	WHO, 2009				
Out-of-pocket expenditure as percentage of private expenditure on health	100.0	WHO, 2009				
Private expenditure on health as percentage of total expenditure on health	18.1	WHO, 2009				
General government expenditure on health as percentage of total government expenditure	13.3	WHO, 2009				

<b>Health Workforce</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Number of nursing and midwifery personnel	545	WHO, 2007				
Nursing and midwifery personnel density (per 10,000 population)	3.2	WHO, 2007				
Number of physicians	52	WHO, 2007				
Physician density (per 10 000 population)	0.23	WHO, 2007				
Number of obstetricians						
Number of paediatricians						
Number of paediatric surgeons						
Number of paediatric cardiac surgeons						
Number of paediatric neurosurgeons						
Number of clinical geneticists						
Number of genetic counsellors						
Number of community health workers						
Number of skilled birth attendants (SBA)						
Density of SBA						
Number of lab staff providing cytogenetic testing						
Number of lab staff providing molecular genetics						
Number of lab staff providing biochemical tests for genetics						

Number of skilled health attendants						
<b>Infrastructure</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Number of maternity units						
Number of services providing specialised care for people with CD						
Number of family planning services						
Number of preconception services						
Number of services providing prenatal care						
Number of services providing newborn care						
Number of facilities providing genetic services						
Number of laboratories providing cytogenetics						
Number of laboratories providing molecular genetics						
Number of laboratories providing biochemical tests for genetics						
Number of facilities for safe terminations of pregnancies for fetal defects						

PPP = purchasing power parity

GDP = gross domestic product

SBA = skilled birth attendant

CD = congenital disorders

**Bhutan****Down's Syndrome****DOWNS Epidemiology 1.1: Country epidemiology**

Epidemiological indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
<b>Year of estimate</b>						
<b>Prevalence at birth and by age-group (/1000)</b>						
Live birth prevalence (LB)			1.70			
Stillbirth prevalence (SB)			0.09			
Total birth prevalence (LB+SB)			1.79			
All age groups						
<1 year olds						
1-4 year olds						
5-14 year olds						
15-44 year olds						
45+ year olds						
<b>Number of cases by age group</b>						
Annual live births			109			
All age groups						
<1 year olds						
1-4 year olds						
5-14 year olds						
15-44 year olds						
45+ year olds						
<b>No. of cases by level of impairment</b>						
No or minor disability						
Moderate disability*						
Severe disability*						
<b>Mortality and morbidity</b>						
Mean life expectancy (yrs)			13			
No. deaths < 1yr			49			
No. deaths 1-4 yrs			17			
No. deaths < 5 yrs			65			
Infant mortality / 1000 LB						
Under-5 mortality / 1000 LB						
Years of life lost			7261			

LB = live births

\* For Down's, Moderate = mild mental retardation and Severe = moderate or worse mental retardation (see Background)

**Bhutan****Down's Syndrome****DOWNS Epidemiology 1.2: International comparison**

Epidemiological indicator	Your chosen estimates	Comparison		
		Country	Region	World
Prevalence at birth and by age-group (/1000 people)		(Asia, South)		
Live birth prevalence (LB)		1.70	1.41	1.52
Stillbirth prevalence (SB)		0.09	0.08	0.09
Total birth prevalence (LB+SB)		1.79	1.49	1.61
All age groups				
<1 year olds				
1-4 year olds				
5-14 year olds				
15-44 year olds				
45+ year olds				
Number of cases by age-group				
Annual live births		109	51739	202668
All age groups			321650	2703100
<1 year olds				
1-4 year olds			102061	480210
5-14 year olds			137963	734321
15-44 year olds			70175	1149318
45+ year olds			11451	339252
No. cases by level of impairment				
No or minor disability				
Moderate disability*				
Severe disability*				
Mortality and morbidity				
Mean life expectancy (yrs)		13	6	14
No. deaths < 1yr		49	25,448	84,981
No. deaths 1-4 yrs		17	9,995	32,644
No. deaths < 5 yrs		65	35,443	117,624
Infant mortality / 1000 LB				
Under-5 mortality / 1000 LB				
Years of life lost		7261	3,824,492	13,333,159

LB = live births

\* For Down's, Moderate = mild mental retardation and Severe = moderate or worse mental retardation (see Background)

**Bhutan****Down's Syndrome****DOWNS Epidemiology 2.1: Data on affected pregnancies: Research studies**

Study author, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies listed above (or in section DOWNS-E2.1 of the Tool), enter the best estimates for the prevalence of affected births and terminations in the country, and a range of values to reflect uncertainty or within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	LB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			
Estimates for the total country/territory	Number of affected still births	SB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			
Estimates for the total country/territory	Number of terminations of pregnancy due to condition	ToP / 1000 women aged 15-44	Comments
Best estimate			
Lower estimate			
Higher estimate			

TB = total births (live births + stillbirths); ToP = termination of pregnancy



**Bhutan****Down's Syndrome****DOWNS Epidemiology 2.2: Data on affected pregnancies: Surveillance**

Based on surveillance data, enter the best estimates for the prevalence of the condition in live births, still births and pregnancy terminations in the country. Give a range of values to reflect uncertainty and within-country variation, and use comments for information on data quality, uncertainty and representativeness.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	Birth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

Estimates for the total country/territory	Number of affected still births	Stillbirth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

Estimates for the total country/territory	Number of ToP due to condition	ToP / 1000 women aged 15-44	Comments
Best estimate			
Lower estimate			
Higher estimate			

TB = total births (live births + stillbirths); ToP = termination of pregnancy

**Bhutan****Down's Syndrome****DOWNS Epidemiology 2.3: Data on affected pregnancies: Other sources**

	Source 1:	Source 2:	Notes
Enter year and source of data – use last year with information available.			
<b>Basic Numbers</b>			
Number of affected live births / year, from data source			
Total number of live births / year, from data source			
Number of affected still births / year, from data source			
Total number of still births / year, from data source			
Number of ToP for affected fetus / year from data source			
Total number of affected births / year (live and still)	0	0	Number of affected live births + Number of affected still births
Total number of births / year, from data source	0	0	Total number of live births + Total number of still births
Total number of ToP / year, from data source			
Total number of women aged 15-44			
<b>Live birth prevalence: recorded and estimated</b>			
Recorded live birth prevalence (affected recorded live births / 1000 total births)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected live births in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded live births (number of recorded live births / total live births in country or territory)			Range: 0 to 1
Estimated live birth prevalence (recorded prevalence / completeness)	#DIV/0!	#DIV/0!	
Estimated true number of affected live births in data source (number of recorded affected live births / completeness)	#DIV/0!	#DIV/0!	
Estimated number of affected live births in total population (number of affected live births from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	
<b>Stillbirth prevalence: recorded and estimated</b>			
Recorded stillbirth prevalence (affected recorded still births / 1000 recorded total births)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected stillbirths in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded stillbirths (number of recorded still births / total still births in country or territory)			Range: 0 to 1
Estimated stillbirth prevalence (recorded prevalence / completeness)	#DIV/0!	#DIV/0!	
Estimated true number of affected stillbirths in data source (number of recorded affected still births / completeness)	#DIV/0!	#DIV/0!	

Estimated number of affected stillbirths in total population (number of affected stillbirths from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	
--	---------	---------	--

<b>ToP prevalence: recorded and estimated</b>			
Recorded ToP prevalence (ToP in affected fetuses / 1000 women aged 15-44)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected pregnancy terminations in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded ToP (number of recorded ToP / total ToP in country or territory)			Range: 0 to 1
Estimated ToP prevalence (recorded prevalence / estimated completeness)	#DIV/0!	#DIV/0!	
Estimated true number of ToP in data source (number of recorded ToP / completeness)	#DIV/0!	#DIV/0!	
Estimated number of ToP in total population (number of ToP from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	
<b>Based on the sources above</b> , enter the best prevalence estimates for your population, and a range of values to reflect uncertainty of estimates and within country variation.			
If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).			
<b>Estimates for the whole country/territory</b>	<b>Number of affected live births</b>	<b>LB prevalence / 1000 TB</b>	
Best estimate			
Lower estimate			
Higher estimate			
<b>Estimates for the whole country/territory</b>	<b>Number of affected stillbirths</b>	<b>SB prevalence / 1000 TB</b>	
Best estimate			
Lower estimate			
Higher estimate			
<b>Estimates for the whole country/territory</b>	<b>Number of ToP due to condition</b>	<b>ToP /1000 women aged 15-44</b>	
Best estimate			
Lower estimate			
Higher estimate			
TB = total births (live births + stillbirths); ToP = termination of pregnancy			

**Bhutan****Down's Syndrome****DOWNS Epidemiology 2.4: Summary of affected pregnancies**

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Number of annual affected <b>live births</b>			109			
Annual birth prevalence / 1000 TB			1.70			
Number of annual affected <b>stillbirths</b>			6			
Annual still birth prevalence / 1000 TB			0.09			
Number of <b>terminations of pregnancy</b> in affected fetuses/ year						
Affected ToP / 1000 women 15-44/ year						

If there are specific sub-types of condition, you can repeat this exercise below. However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

TB = total births (live births + stillbirths); ToP = termination of pregnancy

**Bhutan****Down's Syndrome****DOWNS Epidemiology 2.5: Sub-population variation in affected pregnancies**

If the birth prevalence rates vary by population sub-group (e.g. geographically or by another factor), indicate any population groups with different prevalence estimates from the whole population and describe reasons for variation. If a group is substantially different from the general population, you may wish to conduct a needs assessment for that group alone.

Population sub-group	Number of affected live births	LB prevalence / 1000 TB	Reason for variation

Population sub-group	Number of affected stillbirths	SB prevalence / 1000 TB	Reason for variation

Population sub-group	Number of ToP in affected pregnancies	ToP prevalence / 1000 women aged 15-44	Reason for variation

TB = total births (live births + stillbirths); ToP = termination of pregnancy

**Bhutan****Down's Syndrome****DOWNS Epidemiology 3.1: Mortality data: Research studies**

Source, year, site	Sample size	Age group	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for the specific mortality by age-group e.g. infant, under 5s, etc, as appropriate, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Mortality estimates	Number of deaths	Ratio (deaths / 1000 LB)	Comments
<b>Neonatal group (&lt;28 days)</b>			
Best estimate			
Lower estimate			
Higher estimate			
<b>Infant group (&lt;1 year)</b>			
Best estimate			
Lower estimate			
Higher estimate			
<b>Under-5 group (&lt;5 years)</b>			
Best estimate			
Lower estimate			
Higher estimate			
<b>Other age group:</b>			
Best estimate			
Lower estimate			
Higher estimate			

LB = live births

**Bhutan****Down's Syndrome****DOWNS Epidemiology 3.2: Mortality data: Vital registration data**

Fill in the blank cells based on your vital registration data.	
Enter year and source of data	
<b>Registered data</b>	
Total registered live births	
Registered condition-specific neonatal deaths (first 28 days of life)	
Registered condition-specific infant deaths (first year of life)	
Registered condition-specific under-5 deaths (first 5 years of life)	
Registered condition-specific neonatal mortality ratio (condition-specific neonatal deaths / (Total registered live births / 1000))	#DIV/0!
Registered condition-specific infant mortality ((condition-specific infant deaths / (Total registered live births / 1000))	#DIV/0!
Registered condition-specific under-5 mortality (condition-specific under-5 deaths / (Total registered live births / 1000))	#DIV/0!

**Adjustment for under-ascertainment of cause of death and sub-registration of deaths:** Enter estimates in the highlighted cells. It is not always possible to adjust the estimates, in which case you may give the value '1', accepting that the estimates in these cases will usually be biased towards low values. (Or you may move to the next section.)

It is assumed that under-ascertainment is stable across age-groups; if ascertainment varies by age-group, you could use separate estimates for each age group.

Estimated completeness of recording: what proportion of deaths in affected persons were registered as such?		Range: 0 to 1
Population coverage: what proportion of the total country/territory population is covered by the vital registration?		Range: 0 to 1
Death ascertainment (population coverage x completeness)	0	
<b>Estimated values for the total country/ territory population</b>		
Estimated number of live births in total population (Total registered live births/population coverage)	#DIV/0!	
Estimated number of neonatal deaths in total population (number of deaths registered in neonatal period / ascertainment)	#DIV/0!	
Estimated number of infant deaths in total population (number of deaths registered in first year of life / ascertainment)	#DIV/0!	
Estimated number of under-5 deaths in total population (number of deaths registered in under-5s / ascertainment)	#DIV/0!	
Estimated neonatal mortality ratio (estimated neonatal deaths / 1000 live births)	#DIV/0!	
Estimated infant mortality ratio (estimated infant deaths / 1000 live births)	#DIV/0!	
Estimated under-5 mortality ratio (estimated under-5 deaths / 1000 live births)	#DIV/0!	



**Bhutan****Down's Syndrome****DOWNS Epidemiology 3.3: Mortality data: Other sources**

Source, year, site	Sample size	Age group	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Neonatal mortality		Infant mortality		Under-5 mortality	
Estimates for the total country/territory	Value	Ratio/1000 LB	Value	Ratio/1000 LB	Value	Ratio/1000 LB
Best estimate						
Lower estimate						
Higher estimate						

**Bhutan****Down's Syndrome****DOWNS Epidemiology 3.4: Summary mortality estimates**

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Year of data collection						
Number of annual deaths in affected persons						
Number of annual live births (in 1000s)			64			
Number of annual affected neonatal deaths						
Number of affected neonatal deaths / 1000 LB						
Number of annual affected infant deaths			49			
Number of affected infant deaths/ 1000 LB						
Number of annual affected under-5 deaths			65			
Number of affected under-5 deaths / 1000 LB						
Mean life expectancy at birth in affected people			13			
Other indicators (e.g. survival following surgical procedure, etc)						

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Bhutan****Down's Syndrome****DOWNS Epidemiology 3.5: Sub-population variation in mortality**

Age group: neonatal Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific neonatal mortality ratio / 1000 LB	Reason for variation

Age group: infant Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific infant mortality ratio / 1000 LB	Reason for variation

Age group: under 5 Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific under-5 mortality ratio / 1000 LB	Reason for variation

Age group: ..... Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific mortality ratio / 1000 population	Reason for variation

**Bhutan****Down's Syndrome****DOWNS Epidemiology 4.1: Population prevalence: Research studies**

Study, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for population prevalence, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Bhutan****Down's Syndrome****DOWNS Epidemiology 4.2: Population prevalence: Other sources**

Source, year, site	Sample size	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Bhutan****Down's Syndrome****DOWNS Epidemiology 4.3: Population prevalence summary**

Source of estimates	Estimated total population number of affected persons	Range	Estimated total population prevalence / 1000 persons	Range
1				
2				
3				
4				
5				
PHGDB				
<b>Chosen estimates</b>				

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Bhutan****Down's Syndrome****DOWNS Epidemiology 4.4: Sub-population prevalence variation**

Population sub-group	Number of affected people	Total number of people in population sub-group	Population prevalence per 1000 people	Reason for variation
			#DIV/0!	
			#DIV/0!	
			#DIV/0!	
			#DIV/0!	

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Formula in column D: Number of affected people/ (Total number of people in population subgroup/1000)

**Bhutan****Down's Syndrome****DOWNS Intervention 1: Effects of maternal age**

If you have an estimate for the birth prevalence of chromosomal disorders, you can use the Calculator on the left.

If you have an estimate of the proportion of births that are to mothers aged over 35, you can use the Calculator on the right.

Birth prevalence per 1000 TB		
Proportional birth prevalence due to high maternal age <sup>1</sup>	#DIV/0!	Requires a birth prevalence above
Birth prevalence attributable to high maternal age, per 1000 TB <sup>2</sup>	-0.86	
Baseline prevalence without maternal age effect	0.86	This figure is set at 0.86

TB = total births (live births + stillbirths)

<sup>1</sup>(Birth prevalence – 0.86)/Birth prevalence

<sup>2</sup>Birth prevalence – Baseline prevalence

Proportion of mothers aged >35		Range: 0 to 1
Estimated birth prevalence per 1000 TB <sup>3</sup>	0.86	
Proportional birth prevalence due to high maternal age <sup>4</sup>	0.00	
Birth prevalence attributable to high maternal age, per 1000 TB <sup>5</sup>	0	
Baseline prevalence without maternal age effect	0.86	This figure is set at 0.86

<sup>3</sup>0.86+(7\*Proportion of mothers aged >35)

<sup>4</sup>(Estimated birth prevalence- Baseline prevalence)/Estimated birth prevalence

<sup>5</sup>Estimated birth prevalence\*Proportional birth prevalence



**Bhutan****Down's Syndrome****DOWNS Intervention 2: Effect of prenatal screening and diagnosis and pregnancy termination**

Note: this makes the simplifying assumption that stillbirth is equally likely in cases that are diagnosed as in cases that are not diagnosed.

Assumption: prenatal services are equally used for cases which would lead to stillbirths and live births.

This could overestimate the impact of ToP if in fact ToP is more likely for severe cases that would result in stillbirth.

Conversely, the impact of ToP could be underestimated if screening is only available to high-income women at lower risk.

100% specificity of prenatal diagnosis assumed.

Birth prevalence (allowing for maternal age effect), per 1000 TB		Can be found in sheet DOWNS-Interv1
<b>Variables</b>		
Coverage of prenatal screening and diagnosis		Range: 0 to 1
Proportion of cases diagnosed		Range: 0 to 1
Proportion of diagnosed cases ending in ToP		Range: 0 to 1
<b>Results</b>		
% prevalence reduction due to PNS & ToP <sup>1</sup>	0%	
Prevalence reduction due to PNS & ToP, per 1000 TB <sup>2</sup>	0.000	
Final prevalence: affected live births after PNS & ToP, per 1000 TB <sup>3</sup>	0.000	

PNS = prenatal screening

TB = total births (live births + still births)

ToP = termination of pregnancy

<sup>1</sup>Coverage of PNS and diagnosis X Proportion of cases diagnosed x Proportion of cases ending in ToP

<sup>2</sup>% prevalence reduction due to PNS and ToP x Birth prevalence

<sup>3</sup>Birth prevalence – Prevalence reduction due to PNS & ToP

**Bhutan****Down's Syndrome****DOWNS Needs assessment 1: Quantitative baseline****Table DOWNS-NA1a Burden of Down's Syndrome in pregnancy, at birth and at population level**

Indicator	Chosen estimates			Notes
	Number (n)	n/1000 pregnancy outcomes	Range of prevalence (/1000 TB)	
Annual affected live births (LB)	0	0	0	Drawn from sheet E2.4
Annual affected stillbirths (SB)	0	0	0	Drawn from sheet E2.4
Annual affected births (LB+SB)	0	0		Drawn from sheet E2.4
Annual affected persons (all age groups)	0	0	0	Drawn from sheet E1.1

**Table DOWNS-NA1b Down's Syndrome mortality indicators**

Indicator	Chosen estimates			Notes
	Number (n)	n/1000 LB	Range of prevalence (/1000 LB)	
Annual overall mortality	0			Drawn from sheet E3.4
Annual neonatal mortality	0	0	0	Drawn from sheet E3.4
Annual infant mortality	0	0	0	Drawn from sheet E3.4
Annual under-5 mortality	0	0	0	Drawn from sheet E3.4
Mean life expectancy at birth in affected people	0		0	Drawn from sheet E3.4

TB = total births (live births + stillbirths)

LB = live births

**Bhutan****Down's Syndrome****DOWNS Needs assessment 3: Quantitative assessment of interventions**

<b>Table DOWNS-NA3a</b>	<b>Estimated prevalence in the absence of interventions for Down's Syndrome</b>	
Indicator	Number (n)	Prevalence (n/1000)
Potential live births		
Potential stillbirths		

<b>Table DOWNS-NA3b</b>	<b>Current situation in relation to interventions before birth</b>		
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 TB
Effect of family planning, education			
Effect of prenatal screening			
Effect of prenatal diagnosis			
Effect of ToP			
Overall effect			

<b>Table DOWNS-NA3c</b>	<b>Target situation in relation to interventions before birth</b>		
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 TB
Effect of family planning, education			
Effect of prenatal screening			
Effect of prenatal diagnosis			
Effect of ToP			
Overall effect			

<b>Table DOWNS-NA3d</b>	<b>Current situation in relation to interventions after birth</b>		
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 TB
Effect of newborn diagnosis			
Effect of management and treatment			
Effect of social care and support			
Overall effect			

<b>Table DOWNS-NA3e</b>	<b>Target situation in relation to interventions after birth</b>		
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 TB
Effect of newborn diagnosis			
Effect of management and treatment			
Effect of social care and support			
Overall effect			

<b>Table DOWNS-NA3f</b>	<b>Current and desired outcomes</b>			
	<b>Current situation</b>		<b>Target situation</b>	
Indicator	Annual number (n)	Prevalence (n/1000)	Annual number (n)	Prevalence (n/1000)
<b>Estimated affected pregnancies</b>				
Live births (LB)	0	0		
Stillbirths (SB)	0	0		
Total births (LB+SB)	0	0		
<b>Estimated population prevalence</b>				
All age groups				
<b>Estimated mortality / 1000 live births</b>				
Neonatal deaths	0	0		
Infant deaths	0	0		
Under-5 deaths	0	0		