

PHG Needs Assessment Calculator**Congo****Glucose-6-phosphate dehydrogenase deficiency**

Welcome to the PHG Health Needs Assessment Calculator for Glucose-6-phosphate dehydrogenase deficiency. The contents of this file are listed below.

Full name of the sheet	Short name
Country demographic, maternal health and socioeconomic indicators	Demography
Country health service data	HealthServices
G6PD Epidemiology 1.1: Country epidemiology	G6PD-E1.1
G6PD Epidemiology 1.2: International comparison	G6PD-E1.2
G6PD Epidemiology 2.1: Data on affected pregnancies: Research studies	G6PD-E2.1
G6PD Epidemiology 2.2: Data on affected pregnancies: Surveillance	G6PD-E2.2
G6PD Epidemiology 2.3: Data on affected pregnancies: Other sources	G6PD-E2.3
G6PD Epidemiology 2.4: Summary of affected pregnancies	G6PD-E2.4
G6PD Epidemiology 2.5: Sub-population variation in affected pregnancies	G6PD-E2.5
G6PD Epidemiology 3.1: Mortality data: Research studies	G6PD-E3.1
G6PD Epidemiology 3.2: Mortality data: Vital registration data	G6PD-E3.2
G6PD Epidemiology 3.3: Mortality data: Other sources	G6PD-E3.3
G6PD Epidemiology 3.4: Summary mortality estimates	G6PD-E3.4
G6PD Epidemiology 3.5: Sub-population variation in mortality	G6PD-E3.5
G6PD Epidemiology 4.1: Population prevalence: Research studies	G6PD-E4.1
G6PD Epidemiology 4.2: Population prevalence: Other sources	G6PD-E4.2
G6PD Epidemiology 4.3: Summary of population prevalence	G6PD-E4.3
G6PD Epidemiology 4.4: Sub-population prevalence variation	G6PD-E4.4
G6PD Interventions 1: Newborn screening and management	G6PD-Interv1
G6PD Needs Assessment Calculator 1: Quantitative baseline	G6PD-NA1
G6PD Needs Assessment Calculator 3: Quantitative assessment of interventions	G6PD-NA3

(There is no sheet G6PD-NA2.)

Congo**Shared Data****Demographic, maternal health and socio-economic indicators**

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Demography information from that Calculator into here. The information should be the same.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Number of persons by age-group and sex	Estimates			Your estimates			Chosen estimates		
Age group	Male	Female	Total	Male	Female	Total	Male	Female	Total
0-4 years						0			0
5-9 years						0			0
10-14 years						0			0
15-19 years						0			0
20-24 years						0			0
25-29 years						0			0
30-34 years						0			0
35-39 years						0			0
40-44 years						0			0
45-49 years						0			0
50-54 years						0			0
55-59 years						0			0
60-64 years						0			0
65+ years						0			0
Total	0	0		0	0	0	0	0	0
Female population aged 15-44 years		0			-			-	
Data year	reported in								
Source, Year									

Ethnicity. Please enter data for the main ethnic groups if you are working with a population that is different from that of the country.

Ethnic group	Number	% population

Fertility and mortality	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Crude birth rate: live births (LB) / year / 1000 population	35	Unicef, 2007				
Still birth rate (SB): Still births (SB) / year / 1000 total births	26	WHO, 2009				
Total births in 1000s (LB+SB) per year	133	Unicef, 2007				
Infant mortality rate: infant deaths / 1000 LB / year	61	UNICEF				
Under-5 mortality rate: U5 deaths / 1000 LB / year	93	(2010), 2010				
Percentage births in women >35 years		(2011), 2010				
Life expectancy at birth (yrs)	55	WHO, 2009				
% of marriages consanguineous						

Maternal health	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Prenatal visits – at least 1 visit (%)	86	WHO, 2005				
Prenatal visits – at least 4 visits (%)	75	WHO, 2005				
Births attended by skilled health personnel (%)	86	WHO, 2005				
Contraception prevalence rate (%)	44.3	WHO, 2005				
Unmet need for family planning (%)	16.2	WHO, 2005				
Total fertility rate	4.3	WHO, 2009				
% home births						
% births at health care services						
Newborn health	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Number of neonatal examinations by SBA / trained staff						
% neonatal examinations by SBA/ trained staff						

Socio-economic indicators	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Gross national income per capita (PPP int. \$)	2810	WHO, 2008				
% population living on < US\$1 per day	54.1	WHO, 2005				
Birth registration coverage (%)	81	WHO, 2005				
Death registration coverage (%)						

LB = live births

PPP = purchasing power parity

SBA = skilled birth attendant

Congo
Shared Data
Health Services Data

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Health Services information from that Calculator into here. The information should be the same.

This section provides health-service-related information for your country.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Health Expenditure	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Per capita total expenditure on health (PPP int. \$)	126	WHO, 2009				
Total expenditure on health as percentage of GDP	3.0	WHO, 2009				
Per capita government expenditure on health (PPP int. \$)	68	WHO, 2009				
External resources for health as percentage of total expenditure on health	7.2	WHO, 2009				
General government expenditure on health as percentage of total expenditure on health	53.8	WHO, 2009				
Out-of-pocket expenditure as percentage of private expenditure on health	100.0	WHO, 2009				
Private expenditure on health as percentage of total expenditure on health	46.2	WHO, 2009				
General government expenditure on health as percentage of total government expenditure	5.3	WHO, 2009				

Health Workforce	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Number of nursing and midwifery personnel	3492	WHO, 2007				
Nursing and midwifery personnel density (per 10,000 population)	8.2	WHO, 2007				
Number of physicians	401	WHO, 2007				
Physician density (per 10,000 population)	0.95	WHO, 2007				
Number of obstetricians						
Number of paediatricians						
Number of paediatric surgeons						
Number of paediatric cardiac surgeons						
Number of paediatric neurosurgeons						
Number of clinical geneticists						
Number of genetic counsellors						
Number of community health workers						
Number of skilled birth attendants (SBA)						
Density of SBA						
Number of lab staff providing cytogenetic testing						

Number of lab staff providing molecular genetics						
Number of lab staff providing biochemical tests for genetics						
Number of skilled health attendants						

Infrastructure	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Number of maternity units						
Number of services providing specialised care for people with CD						
Number of family planning services						
Number of preconception services						
Number of services providing prenatal care						
Number of services providing newborn care						
Number of facilities providing genetic services						
Number of laboratories providing cytogenetics						
Number of laboratories providing molecular genetics						
Number of laboratories providing biochemical tests for genetics						
Number of facilities for safe terminations of pregnancies for fetal defects						

PPP = purchasing power parity

GDP = gross domestic product

SBA = skilled birth attendant

CD = congenital disorders

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 1.1: Country epidemiology**

Epidemiological indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Year of estimate						
Prevalence at birth and by age-group (/1000)						
Live birth prevalence (LB)			1.74			
Stillbirth prevalence (SB)			0.00			
Total birth prevalence (LB+SB)			1.74			
All age groups						
<1 year olds			1.45			
1-4 year olds						
5-14 year olds						
15-44 year olds						
45+ year olds						
Number of cases by age group						
Annual live births			300			
All age groups						
<1 year olds			190			
1-4 year olds						
5-14 year olds						
15-44 year olds						
45+ year olds						
No. of cases by level of impairment						
No or minor disability						
Moderate disability			95			
Severe disability*			95			
Mortality and morbidity						
Mean life expectancy (yrs)						
No. deaths < 1yr			109			
No. deaths 1-4 yrs			0			
No. deaths < 5 yrs			109			
Infant mortality / 1000 LB			0.64			
Under-5 mortality / 1000 LB			0.64			
Years of life lost						

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 1.2: International comparison**

Epidemiological indicator	Your chosen estimates	Comparison		
		Country	Region	World
Prevalence at birth and by age-group (/1000 people)		(Sub-Saharan Africa, Central)		
Live birth prevalence (LB)		1.74	1.71	0.55
Stillbirth prevalence (SB)		0.00	0.00	0.00
Total birth prevalence (LB+SB)		1.74	1.71	0.55
All age groups				
<1 year olds		1.45		
1-4 year olds				
5-14 year olds				
15-44 year olds				
45+ year olds				
Number of cases by age-group				
Annual live births		300	6721	72687
All age groups				
<1 year olds		190	3454	41883
1-4 year olds				
5-14 year olds				
15-44 year olds				
45+ year olds				
No. cases by level of impairment				
No or minor disability				
Moderate disability		95	1727	20942
Severe disability		95	5116	1455
Mortality and morbidity				
Mean life expectancy (yrs)				
No. deaths < 1yr		109	3242	30779
No. deaths 1-4 yrs		0	0	0
No. deaths < 5 yrs		109	3242	30779
Infant mortality / 1000 LB		0.64	0.83	0.23
Under-5 mortality / 1000 LB		0.64	0.83	0.23
Years of life lost				

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 2.1: Data on affected pregnancies: Research studies**

Study author, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies listed above (or in section SCD-E2.1 of the Tool), enter the best estimates for the prevalence of affected births and terminations in the country, and a range of values to reflect uncertainty or within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	LB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			
Estimates for the total country/territory	Number of affected stillbirths	SB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

TB = total births (live births + stillbirths)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 2.2: Data on affected pregnancies: Surveillance**

Based on surveillance data, enter the best estimates for the prevalence of the condition in live births and still births in the country. Give a range of values to reflect uncertainty and within-country variation, and use comments for information on data quality, uncertainty and representativeness.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	Birth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

Estimates for the total country/territory	Number of affected stillbirths	Stillbirth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

TB = total births (live births + stillbirths)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 2.3: Data on affected pregnancies: Other sources**

	Source 1:	Source 2:	Notes
Enter year and source of data – use last year with information available.			
Basic Numbers			
Number of affected live births / year, from data source			
Total number of live births / year, from data source			
Number of affected still births / year, from data source			
Total number of stillbirths / year, from data source			
Total number of affected births / year (live and still)	0	0	
Total number of births / year, from data source	0	0	
Total number of women aged 15-44			
Live birth prevalence: recorded and estimated			
Recorded live birth prevalence (affected recorded live births / 1000 total births)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected live births in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded live births (number of recorded live births / total live births in country or territory)			Range: 0 to 1
Estimated live birth prevalence (recorded prevalence / completeness)	#DIV/0!	#DIV/0!	
Estimated true number of affected live births in data source (number of recorded affected live births / completeness)	#DIV/0!	#DIV/0!	
Estimated number of affected live births in total population (number of affected live births from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	
Stillbirth prevalence: recorded and estimated			
Recorded stillbirth prevalence (affected recorded still births / 1000 recorded total births)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected stillbirths in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded stillbirths (number of recorded still births / total still births in country or territory)			Range: 0 to 1
Estimated stillbirth prevalence (recorded prevalence / completeness)	#DIV/0!	#DIV/0!	
Estimated true number of affected stillbirths in data source (number of recorded affected still births / completeness)	#DIV/0!	#DIV/0!	
Estimated number of affected still births in total population (number of affected still births from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	

Based on the sources above, enter the best prevalence estimates for your population, and a range of values to reflect uncertainty of estimates and within country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the whole country/territory	Number of affected live births	LB prevalence / 1000 TB
Best estimate		
Lower estimate		
Higher estimate		
Estimates for the whole country/territory	Number of affected stillbirths	SB prevalence / 1000 TB
Best estimate		
Lower estimate		
Higher estimate		

TB = total births (live births + stillbirths)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 2.4: Summary of affected pregnancies**

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Number of annual affected live births			300			
Annual birth prevalence / 1000 TB			1.74			
Number of annual affected still births			0			
Stillbirth prevalence / 1000 TB/year			0.00			

If there are specific sub-types of condition, you can repeat this exercise below. However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

TB = total births (live births + stillbirths)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 2.5: Sub-population variation in affected pregnancies**

If the birth prevalence rates vary by population sub-group (e.g. geographically or by another factor), indicate any population groups with different prevalence estimates from the whole population and describe reasons for variation. If a group is substantially different from the general population, you may wish to conduct a needs assessment for that group alone.

Population sub-group	Number of affected live births	LB prevalence / 1000 TB	Reason for variation

Population sub-group	Number of affected stillbirths	SB prevalence / 1000 TB	Reason for variation

TB = total births (live births + stillbirths)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 3.1: Mortality data: Research studies**

Source, year, site	Sample size	Age group	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for the specific mortality by age-group e.g. infant, under-5s, etc., as appropriate, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Mortality estimates	Number of deaths	Ratio (deaths / 1000 LB)	Comments
Neonatal group (<28 days)			
Best estimate			
Lower estimate			
Higher estimate			
Infant group (<1 year)			
Best estimate			
Lower estimate			
Higher estimate			
Under-5 group (<5 years)			
Best estimate			
Lower estimate			
Higher estimate			
Other age group:			
Best estimate			
Lower estimate			
Higher estimate			

LB = live births

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 3.2: Mortality data: Vital registration data**

Fill in the blank cells based on your vital registration data.	
Enter year and source of data	
Registered data	
Total registered live births	
Registered condition-specific neonatal deaths (first 28 days of life)	
Registered condition-specific infant deaths (first year of life)	
Registered condition-specific under-5 deaths (first 5 years of life)	
Registered condition-specific neonatal mortality ratio (condition-specific neonatal deaths / 1000 live births in the same year)	#DIV/0!
Registered condition-specific infant mortality (condition-specific infant deaths / 1000 live births in the same year)	#DIV/0!
Registered condition-specific under-5 mortality (condition-specific under-5 deaths / 1000 live births in the same year)	#DIV/0!

Adjustment for under-ascertainment of cause of death and sub-registration of deaths: Enter estimates in the highlighted cells. It is not always possible to adjust the estimates, in which case you may give the value '1', accepting that the estimates in these cases will usually be biased towards low values. (Or you may move to the next section.)

It is assumed that under-ascertainment is stable across age-groups; if ascertainment varies by age-group, you could use separate estimates for each age group.

Estimated completeness of recording: what proportion of deaths in affected persons were registered as such?		Range: 0 to 1
Population coverage: what proportion of the total country/territory population is covered by the vital registration?		Range: 0 to 1
Death ascertainment (population coverage x completeness)	0	
Estimated values for the total country/ territory population		
Estimated number of live births in total population	#DIV/0!	
Estimated number of neonatal deaths in total population (number of deaths registered in neonatal period / ascertainment)	#DIV/0!	
Estimated number of infant deaths in total population (number of deaths registered in first year of life / ascertainment)	#DIV/0!	
Estimated number of under-5 deaths in total population (number of deaths registered in under-5s / ascertainment)	#DIV/0!	
Estimated neonatal mortality ratio (estimated neonatal deaths / 1000 live births)	#DIV/0!	
Estimated infant mortality ratio (estimated infant deaths / 1000 live births)	#DIV/0!	
Estimated under-5 mortality ratio (estimated under-5 deaths / 1000 live births)	#DIV/0!	

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 3.3: Mortality data: Other sources**

Source, year, site	Sample size	Age group	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Neonatal mortality		Infant mortality		Under-5 mortality	
Estimates for the total country/territory	Value	Ratio/1000 LB	Value	Ratio/1000 LB	Value	Ratio/1000 LB
Best estimate						
Lower estimate						
Higher estimate						

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 3.4: Summary mortality estimates**

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Year of data collection						
Number of annual deaths in affected persons						
Number of annual live births (in 1000s)			172			
Number of annual affected neonatal deaths			0			
Number of affected neonatal deaths / 1000 LB			0.00			
Number of annual affected infant deaths			109			
Number of affected infant deaths / 1000 LB			0.64			
Number of annual affected under-5 deaths			109			
Number of affected under-5 deaths / 1000 LB			0.64			
Mean life expectancy at birth in affected people						
Other indicators (e.g. survival following surgical procedure, etc)						

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 3.5: Sub-population variation in mortality**

Age group: neonatal Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific neonatal mortality ratio / 1000 LB	Reason for variation

Age group: infant Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific infant mortality ratio / 1000 LB	Reason for variation

Age group: under 5 Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific under-5 mortality ratio / 1000 LB	Reason for variation

Age group: Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific mortality ratio / 1000 population	Reason for variation

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 4.1: Population prevalence: Research studies**

Study, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for population prevalence, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 4.2: Population prevalence: Other sources**

Source, year, site	Sample size	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 4.3: Summary of population prevalence**

Source of estimates	Estimated total population number of affected persons	Range	Estimated total population prevalence / 1000 persons	Range
1				
2				
3				
4				
5				
PHGDB				
Chosen estimates				

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Epidemiology 4.4: Sub-population prevalence variation**

Population sub-group	Number of affected people	Total number of people in population sub-group	Population prevalence per 1000 people	Reason for variation
			#DIV/0!	
			#DIV/0!	
			#DIV/0!	
			#DIV/0!	

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Formula in column D: Number of affected people/ (Total number of people in population subgroup/1000)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Intervention 1: Effects of NBS and treatment on G6PDD****Screened**

Baseline birth prevalence of G6PDD, per 1000 TB*		
Variables		
Coverage of newborn screening		0 Range: 0 to 1
Proportion of positive-screened patients receiving treatment		Range: 0 to 1
Effectiveness of treatment in screened patients		Range: 0 to 1
Results		
Proportional reduction of unmanaged cases of G6PDD through NBS and treatment ¹		0
Prevalence of unmanaged G6PDD due to newborn screening and treatment, per 1000 total births ²		0
Unscreened		
Birth prevalence of unscreened G6PDD, per 1000		0
Variables		
Estimated coverage of clinical diagnosis in unscreened patients		Range: 0 to 1
Proportion of unscreened patients diagnosed clinically receiving treatment		Range: 0 to 1
Effectiveness of treatment in unscreened patients		Range: 0 to 1
Results		
Proportional reduction of unscreened unmanaged cases of G6PDD through clinical diagnosis and treatment ³		0
Prevalence of unscreened unmanaged G6PDD following clinical diagnosis, per 1000 total births ⁴		0

Total

Prevalence of unmanaged G6PDD following treatment in both screen-detected and clinically-detected patients, per 1000 total births ⁵	0
--	---

TB = total births (live births + still births)

G6PDD = Glucose-6-Phosphate Dehydrogenase Deficiency

* If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population)

¹Coverage of newborn screening x Proportion of positive screened patients receiving treatment x Effectiveness of treatment in screened patients

²Baseline birth prevalence -(proportional reduction of unmanaged cases of G6PDD through NBS and treatment x Baseline birth prevalence)

³Coverage of clinical diagnosis in unscreened patients x Proportion of clinically diagnosed patients receiving treatment x Effectiveness of treatment in unscreened patients

⁴Birth prevalence of unscreened G6PDD – (Proportional reduction of unscreened unmanaged cases through clinical diagnosis x Birth prevalence of unscreened G6PDD)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Needs Assessment Calculator 1: Quantitative baseline****Table SCD-NA1a Burden of Glucose-6-phosphate dehydrogenase deficiency in pregnancy, at birth and at population level**

Indicator	Chosen estimates			Notes
	Number (n)	n/1000 TB	Range of prevalence (/1000 TB)	
Annual affected live births (LB)	0	0	0	Drawn from sheet E2.4
Annual affected stillbirths (SB)	0	0	0	Drawn from sheet E2.4
Annual affected births (LB+SB)	0	0		Drawn from sheet E2.4
Annual affected persons (all age groups)	0	0	0	Drawn from sheet E1.1

Table SCD-NA1b Glucose-6-phosphate dehydrogenase deficiency mortality indicators

Indicator	Chosen estimates			Notes
	Number (n)	n/1000 LB	Range of prevalence (/1000 TB)	
Annual overall mortality	0			Drawn from sheet E3.4
Annual neonatal mortality	0	0	0	Drawn from sheet E3.4
Annual infant mortality	0	0	0	Drawn from sheet E3.4
Annual under-5 mortality	0	0	0	Drawn from sheet E3.4
Mean life expectancy at birth among affected people	0		0	Drawn from sheet E3.4

TB = total births (live births + stillbirths)

Congo**Glucose-6-phosphate dehydrogenase deficiency****G6PDD Needs Assessment Calculator 3: Quantitative assessment of interventions**

Table G6PDD-NA3a	Estimated prevalence in the absence of interventions for Glucose-6-phosphate dehydrogenase deficiency	
Indicator	Number (n)	Prevalence (n/1000)
Potential live births		
Potential still births		

Table G6PDD-NA3b	Current situation in relation to interventions before birth		
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 LB
Effect of family planning, education			
Effect of population carrier screening			
Effect of preconception screening			
Effect of prenatal screening			
Overall effect			

Table G6PDD-NA3c	Target situation in relation to interventions before birth		
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 LB
Effect of family planning, education			
Effect of population carrier screening			
Effect of preconception screening			
Effect of prenatal screening			
Overall effect			

Table G6PDD-NA3d	Current situation in relation to interventions after birth		
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 LB
Effect of newborn screening			
Effect of newborn diagnosis			
Treatment of newborn			
Haemolysis prevention			
Haemolysis treatment post neonatal			
Effect of social care and support			
Overall effect			

Table G6PDD-NA3e	Target situation in relation to interventions after birth		
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 LB
Effect of newborn screening			
Effect of newborn diagnosis			
Treatment of newborn			
Haemolysis prevention			
Haemolysis treatment post neonatal			
Effect of social care and support			
Overall effect			

Table G6PDD-NA3f	Current and desired outcomes			
	Current situation		Target situation	
Indicator	Annual number (n)	Incidence (n/1000)	Annual number (n)	Incidence (n/1000)
Estimated affected pregnancies				
Live births (LB)	0	0		
Still births (SB)	0	0		
All births (LB+SB)	0	0		
Estimated population prevalence				
All age groups				
Estimated mortality				
Neonatal deaths	0	0		
Infant deaths	0	0		
Under-5 deaths	0	0		