

**PHG Needs Assessment Calculator****Nigeria****Rhesus Haemolytic Disease of the Newborn**

Welcome to the PHG Health Needs Assessment Calculator for Rhesus Haemolytic Disease of the Newborn. The contents of this file are listed below.

Full name of the sheet	Short name
Country demographic, maternal health and socioeconomic indicators	Demography
Country health service data	HealthServices
RHD Epidemiology 1.1: Country epidemiology	RHD-E1.1
RHD Epidemiology 1.2: International comparison	RHD-E1.2
RHD Epidemiology 2.1: Data on affected pregnancies: Research studies	RHD-E2.1
RHD Epidemiology 2.2: Data on affected pregnancies: Surveillance	RHD-E2.2
RHD Epidemiology 2.3: Data on affected pregnancies: Other sources	RHD-E2.3
RHD Epidemiology 2.4: Summary of affected pregnancies	RHD-E2.4
RHD Epidemiology 2.5: Sub-population variation in affected pregnancies	RHD-E2.5
RHD Epidemiology 3.1: Mortality data: Research studies	RHD-E3.1
RHD Epidemiology 3.2: Mortality data: Vital registration data	RHD-E3.2
RHD Epidemiology 3.3: Mortality data: Other sources	RHD-E3.3
RHD Epidemiology 3.4: Summary mortality estimates	RHD-E3.4
RHD Epidemiology 3.5: Sub-population variation in mortality	RHD-E3.5
RHD Epidemiology 4.1: Population prevalence: Research studies	RHD-E4.1
RHD Epidemiology 4.2: Population prevalence: Other sources	RHD-E4.2
RHD Epidemiology 4.3: Summary of population prevalence	RHD-E4.3
RHD Epidemiology 4.4: Sub-population prevalence variation	RHD-E4.4
RHD Interventions 1: Prenatal screening and treatment	RHD-Interv1
RHD Interventions 2: Newborn screening and treatment	RHD-Interv2
RHD Needs Assessment Calculator 1: Quantitative baseline	RHD-NA1
RHD Needs Assessment Calculator 3: Quantitative assessment of interventions	RHD-NA3

(There is no sheet RHD-NA2.)

**Nigeria****Shared Data****Demographic, maternal health and socio-economic indicators**

**Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Demography information from that Calculator into here. The information should be the same.**

**By default, the Toolkit contains information at the national level.**

**If you would like to use a different population, then replace country information with that of your specific population of interest.**

Number of persons by age-group and sex	Estimates			Your estimates			Chosen estimates		
Age group	Male	Female	Total	Male	Female	Total	Male	Female	Total
0-4 years	11569218	11025749	22594967			0			0
5-9 years	10388611	9616769	20005380			0			0
10-14 years	8504319	7631631	16135950			0			0
15-19 years	7536532	7362887	14899419			0			0
20-24 years	6237549	7197530	13435079			0			0
25-29 years	5534458	6676968	12211426			0			0
30-34 years	4505186	4962352	9467538			0			0
35-39 years	3661133	3670622	7331755			0			0
40-44 years	3395489	3060981	6456470			0			0
45-49 years	2561526	2029767	4591293			0			0
50-54 years	2363937	1885282	4249219			0			0
55-59 years	1189770	876477	2066247			0			0
60-64 years	1363219	1087067	2450286			0			0
65+ years	2534541	2002220	4536761			0			0
Total	0	0	1.4E+008	0	0	0	0	0	0
Female population aged 15-44 years		0			-			-	
Data year	2006 reported in 2010								
Source, Year	UN 2011								

**Ethnicity. Please enter data for the main ethnic groups if you are working with a population that is different from that of the country.**

Ethnic group	Number	% population

<b>Fertility and mortality</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Crude birth rate: live births (LB) / year / 1000 population	40	Unicef, 2007				
Still birth rate (SB): Still births (SB) / year / 1000 total births	42	WHO, 2009				
Total births in 1000s (LB+SB) per year	5959	Unicef, 2007				
Infant mortality rate: infant deaths / 1000 LB / year	88	UNICEF				
Under-5 mortality rate: U5 deaths / 1000 LB / year	143	UNICEF, 2010				
Percentage births in women >35 years		(2011), 2010				
Life expectancy at birth (yrs)	54	WHO, 2009				
% of marriages consanguineous						

<b>Maternal health</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Prenatal visits – at least 1 visit (%)	58	WHO, 2008				
Prenatal visits – at least 4 visits (%)	45	WHO, 2008				
Births attended by skilled health personnel (%)	38.9	WHO, 2008				
Contraception prevalence rate (%)	14.6	WHO, 2008				
Unmet need for family planning (%)	20.2	WHO, 2008				
Total fertility rate	5.2	WHO, 2009				
% home births						
% births at health care services						
<b>Newborn health</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Number of neonatal examinations by SBA / trained staff						
% neonatal examinations by SBA/ trained staff						

<b>Socio-economic indicators</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Gross national income per capita (PPP int. \$)	1980	WHO, 2008				
% population living on < US\$1 per day	64.4	WHO, 2004				
Birth registration coverage (%)	30	WHO, 2008				
Death registration coverage (%)						

LB = live births

PPP = purchasing power parity

SBA = skilled birth attendant

**Nigeria****Shared Data****Health Services Data**

**Please read first!** If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Health Services information from that Calculator into here. The information should be the same.

This section provides health-service-related information for your country.

**By default, the Toolkit contains information at the national level.**

**If you would like to use a different population, then replace country information with that of your specific population of interest.**

<b>Health Expenditure</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Per capita total expenditure on health (PPP int. \$)	136	WHO, 2009				
Total expenditure on health as percentage of GDP	5.8	WHO, 2009				
Per capita government expenditure on health (PPP int. \$)	50	WHO, 2009				
External resources for health as percentage of total expenditure on health	4.9	WHO, 2009				
General government expenditure on health as percentage of total expenditure on health	36.3	WHO, 2009				
Out-of-pocket expenditure as percentage of private expenditure on health	95.6	WHO, 2009				
Private expenditure on health as percentage of total expenditure on health	63.7	WHO, 2009				
General government expenditure on health as percentage of total government expenditure	6.4	WHO, 2009				

<b>Health Workforce</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Number of nursing and midwifery personnel	224943	WHO, 2008				
Nursing and midwifery personnel density (per 10,000 population)	16.1	WHO, 2008				
Number of physicians	55376	WHO, 2008				
Physician density (per 10,000 population)	3.95	WHO, 2008				
Number of obstetricians						
Number of paediatricians						
Number of paediatric surgeons						
Number of paediatric cardiac surgeons						
Number of paediatric neurosurgeons						
Number of clinical geneticists						
Number of genetic counsellors						
Number of community health workers						
Number of skilled birth attendants (SBA)						

Density of SBA						
Number of lab staff providing cytogenetic testing						
Number of lab staff providing molecular genetics						
Number of lab staff providing biochemical tests for genetics						
Number of skilled health attendants						

<b>Infrastructure</b>	<b>Estimate</b>	<b>Source, Year</b>	<b>Your estimate</b>	<b>Source, Year</b>	<b>Chosen estimate</b>	<b>Source, Year</b>
Number of maternity units						
Number of services providing specialised care for people with CD						
Number of family planning services						
Number of preconception services						
Number of services providing prenatal care						
Number of services providing newborn care						
Number of facilities providing genetic services						
Number of laboratories providing cytogenetics						
Number of laboratories providing molecular genetics						
Number of laboratories providing biochemical tests for genetics						
Number of facilities for safe terminations of pregnancies for fetal defects						

PPP = purchasing power parity

GDP = gross domestic product

SBA = skilled birth attendant

CD = congenital disorders

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 1.1: Country epidemiology**

Epidemiological indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
<b>Year of estimate</b>						
<b>Prevalence at birth and by age-group (/1000)</b>						
Live birth prevalence (LB)			0.94			
Stillbirth prevalence (SB)			0.94			
Total birth prevalence (LB+SB)			1.87			
All age groups						
<1 year olds			0.47			
1-4 year olds			0.40			
5-14 year olds			0.00			
15-44 year olds			0.00			
45+ year olds			0.00			
<b>Number of cases by age group</b>						
Annual live births			4,978			
All age groups						
<1 year olds			2,489			
1-4 year olds			2,116			
5-14 year olds			0			
15-44 year olds			0			
45+ year olds			0			
<b>No. of cases by level of impairment</b>						
No or minor disability						
Moderate disability			2,116			
Severe disability			373			
<b>Mortality and morbidity</b>						
Mean life expectancy (yrs)						
No. deaths < 1yr			2,489			
No. deaths 1-4 yrs			373			
No. deaths < 5 yrs			2,862			
Infant mortality / 1000 LB			0.47			
Under-5 mortality / 1000 LB			0.54			
Years of life lost						

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 1.2: International comparison**

	Your chosen estimates	Comparison		
Epidemiological indicator		Country	Region	World
Prevalence at birth and by age-group (/1000 people)		(Sub-Saharan Africa, West)		
Live birth prevalence (LB)		0.94	0.87	0.31
Stillbirth prevalence (SB)		0.94	0.87	0.31
Total birth prevalence (LB+SB)		1.87	1.74	0.63
All age groups				
<1 year olds		0.47		
1-4 year olds		0.40		
5-14 year olds		0.00		
15-44 year olds		0.00		
45+ year olds		0.00		
Number of cases by age-group				
Annual live births		4,978	10195	41746
All age groups				
<1 year olds		2,489	5097	20873
1-4 year olds		2,116	4333	17742
5-14 year olds		0	0	0
15-44 year olds		0	0	0
45+ year olds		0	0	0
No. cases by level of impairment				
No or minor disability			0	0
Moderate disability		2,116	4333	17742
Severe disability		373	218	184
Mortality and morbidity				
Mean life expectancy (yrs)				
No. deaths < 1yr		2,489	5097	20873
No. deaths 1-4 yrs		373	765	3131
No. deaths < 5 yrs		2,862	5862	24004
Infant mortality / 1000 LB		0.47	0.43	0.16
Under-5 mortality / 1000 LB		0.54	0.50	0.18
Years of life lost				

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 2.1: Data on affected pregnancies: Research studies**

Study author, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies listed above (or in section RHD-E2.1 of the Tool), enter the best estimates for the prevalence of affected births and terminations in the country, and a range of values to reflect uncertainty or within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	LB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			
Estimates for the total country/territory	Number of affected stillbirths	SB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

TB = total births (live births + stillbirths);



**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 2.2: Data on affected pregnancies: Surveillance**

Based on surveillance data, enter the best estimates for the prevalence of the condition in live births and still births. Give a range of values to reflect uncertainty and within-country variation, and use comments for information on data quality, uncertainty and representativeness.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	Birth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

Estimates for the total country/territory	Number of affected stillbirths	Stillbirth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

TB = total births (live births + stillbirths)

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 2.3: Data on affected pregnancies: Other sources**

	Source 1:	Source 2:	Notes
Enter year and source of data – use last year with information available.			
<b>Basic Numbers</b>			
Number of affected live births / year, from data source			
Total number of live births / year, from data source			
Number of affected still births / year, from data source			
Total number of stillbirths / year, from data source			
Total number of affected births / year (live and still)	0	0	
Total number of births / year, from data source	0	0	
Total number of women aged 15-44			
<b>Live birth prevalence: recorded and estimated</b>			
Recorded live birth prevalence (affected recorded live births / 1000 recorded total births)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected live births in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded live births (number of recorded live births / total live births in country or territory)			Range: 0 to 1
Estimated live birth prevalence (recorded prevalence / completeness)	#DIV/0!	#DIV/0!	
Estimated true number of affected live births in data source (number of recorded affected live births / completeness)	#DIV/0!	#DIV/0!	
Estimated number of affected live births in total population (number of affected live births from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	
<b>Stillbirth prevalence: recorded and estimated</b>			
Recorded stillbirth prevalence (affected recorded still births / 1000 recorded total births)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected stillbirths in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded stillbirths (number of recorded still births / total still births in country or territory)			Range: 0 to 1
Estimated stillbirth prevalence (recorded prevalence / completeness)	#DIV/0!	#DIV/0!	
Estimated true number of affected stillbirths in data source (number of recorded affected still births / completeness)	#DIV/0!	#DIV/0!	
Estimated number of affected stillbirths in total population (number of affected still births from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	

**Based on the sources above**, enter the best prevalence estimates for your population, and a range of values to reflect uncertainty of estimates and within country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the whole country/territory	Number of affected live births	LB prevalence / 1000 TB
Best estimate		
Lower estimate		
Higher estimate		
Estimates for the whole country/territory	Number of affected still births	SB prevalence / 1000 TB
Best estimate		
Lower estimate		
Higher estimate		

TB = total births (live births + stillbirths)

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 2.4: Summary of affected pregnancies**

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Number of annual affected <b>live births</b>			4,978			
Annual birth prevalence / 1000 TB			0.94			
Number of annual affected <b>still births</b>			4,978			
Stillbirth prevalence / 1000 TB/year			0.94			

If there are specific sub-types of condition, you can repeat this exercise below. However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

TB = total births (live births + stillbirths)

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 2.5: Sub-population variation in affected pregnancies**

If the birth prevalence rates vary by population sub-group (e.g. geographically or by another factor), indicate any population groups with different prevalence estimates from the whole population and describe reasons for variation. If a group is substantially different from the general population, you may wish to conduct a needs assessment for that group alone.

Population sub-group	Number of affected live births	LB prevalence / 1000 TB	Reason for variation

Population sub-group	Number of affected stillbirths	SB prevalence / 1000 TB	Reason for variation

--	--	--	--

TB = total births (live births + stillbirths)

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 3.1: Mortality data: Research studies**

Source, year, site	Sample size	Age group	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for the specific mortality by age-group e.g. infant, under-5s, etc., as appropriate, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Mortality estimates	Number of deaths	Ratio (deaths / 1000 LB)	Comments
<b>Neonatal group (&lt;28 days)</b>			
Best estimate			
Lower estimate			
Higher estimate			
<b>Infant group (&lt;1 year)</b>			
Best estimate			
Lower estimate			
Higher estimate			
<b>Under-5 group (&lt;5 years)</b>			
Best estimate			
Lower estimate			
Higher estimate			
<b>Other age group:</b>			
Best estimate			
Lower estimate			
Higher estimate			

LB = live births

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 3.2: Mortality data: Vital registration data**

Fill in the blank cells based on your vital registration data.	
Enter year and source of data	
<b>Registered data</b>	
Total registered live births	
Registered condition-specific neonatal deaths (first 28 days of life)	
Registered condition-specific infant deaths (first year of life)	
Registered condition-specific under-5 deaths (first 5 years of life)	
Registered condition-specific neonatal mortality ratio (condition-specific neonatal deaths / 1000 live births in the same year)	#DIV/0!
Registered condition-specific infant mortality (condition-specific infant deaths / 1000 live births in the same year)	#DIV/0!
Registered condition-specific under-5 mortality (condition-specific under-5 deaths / 1000 live births in the same year)	#DIV/0!

**Adjustment for under-ascertainment of cause of death and sub-registration of deaths:** Enter estimates in the highlighted cells. It is not always possible to adjust the estimates, in which case you may give the value '1', accepting that the estimates in these cases will usually be biased towards low values. (Or you may move to the next section.)  
It is assumed that under-ascertainment is stable across age-groups; if ascertainment varies by age-group, you could use separate estimates for each age group.

Estimated completeness of recording: what proportion of deaths in affected persons were registered as such?		Range: 0 to 1
Population coverage: what proportion of the total country/territory population is covered by the vital registration?		Range: 0 to 1
Death ascertainment (population coverage x completeness)	0	
<b>Estimated values for the total country/ territory population</b>		
Estimated number of live births in total population	#DIV/0!	
Estimated number of neonatal deaths in total population (number of deaths registered in neonatal period / ascertainment)	#DIV/0!	
Estimated number of infant deaths in total population (number of deaths registered in first year of life / ascertainment)	#DIV/0!	
Estimated number of under-5 deaths in total population (number of deaths registered in under-5s / ascertainment)	#DIV/0!	
Estimated neonatal mortality ratio (estimated neonatal deaths / 1000 live births)	#DIV/0!	
Estimated infant mortality ratio (estimated infant deaths / 1000 live births)	#DIV/0!	
Estimated under-5 mortality ratio (estimated under-5 deaths / 1000 live births)	#DIV/0!	

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 3.3: Mortality data: Other sources**

Source, year, site	Sample size	Age group	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Neonatal mortality		Infant mortality		Under-5 mortality	
Estimates for the total country/territory	Value	Ratio/1000 LB	Value	Ratio/1000 LB	Value	Ratio/1000 LB
Best estimate						
Lower estimate						
Higher estimate						



**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 3.4: Summary mortality estimates**

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Year of data collection						
Number of annual deaths in affected persons			7,342			
Number of annual live births (in 1000s)			5,323			
Number of annual affected neonatal deaths			1,991			
Number of affected neonatal deaths / 1000 LB			0.37			
Number of annual affected infant deaths			2,489			
Number of affected infant deaths / 1000 LB			0.47			
Number of annual affected under-5 deaths			2,862			
Number of affected under-5 deaths / 1000 LB			0.54			
Mean life expectancy at birth in affected people						
Other indicators (e.g. survival following surgical procedure, etc)						

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 3.5: Sub-population variation in mortality**

Age group: neonatal Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific neonatal mortality ratio / 1000 LB	Reason for variation

Age group: infant Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific infant mortality ratio / 1000 LB	Reason for variation

Age group: under 5 Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific under-5 mortality ratio / 1000 LB	Reason for variation

Age group: ..... Population sub-group	Number of deaths in affected persons	Cause-specific, group-specific mortality ratio / 1000 population	Reason for variation

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 4.1: Population prevalence: Research studies**

Study, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for population prevalence, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 4.2: Population prevalence: Other sources**

Source, year, site	Sample size	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 4.3: Summary of population prevalence**

Source of estimates	Estimated total population number of affected persons	Range	Estimated total population prevalence / 1000 persons	Range
1				
2				
3				
4				
5				
PHGDB				
<b>Chosen estimates</b>				

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Epidemiology 4.4: Sub-population prevalence variation**

Population sub-group	Number of affected people	Total number of people in population sub-group	Population prevalence per 1000 people	Reason for variation
			#DIV/0!	
			#DIV/0!	
			#DIV/0!	
			#DIV/0!	

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Formula in column D: Number of affected people/ (Total number of people in population subgroup/1000)

**Nigeria****Rhesus Haemolytic Disease of the Newborn****SCD Intervention 1: Effects of prenatal screening and treatment**

Baseline prevalence of RHD per 1000 TB		
Variables		
Coverage of prenatal screening		Range: 0 to 1
Proportion of RhD negative women receiving anti-D		Range: 0 to 1
Effectiveness of anti-D in RhD negative women		Range: 0 to 1
Results		
% prevalence reduction due to PNS & treatment <sup>1</sup>	0%	
Prevalence reduction due to PNS & treatment, per 1000 TB <sup>2</sup>	0.000	
Final prevalence of RHD-affected live births and still births after PNS & treatment, per 1000 TB <sup>3</sup>	0.000	

RHD = Rhesus Haemolytic Disease of the Newborn

PNS = prenatal screening

TB = total births (live births + still births)

<sup>1</sup>(Coverage of PNS X Proportion of women receiving anti-D) X Effectiveness of anti-D

<sup>2</sup>% prevalence reduction due to PNS and treatment X Baseline prevalence of RHD

<sup>3</sup>Baseline prevalence of RHD – Prevalence reduction due to PNS and treatment

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Intervention 2: Effects of NBS and management on Rhesus Haemolytic Disease of the Newborn**

Baseline birth prevalence of RHD, per 1000 LB		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of positive-screened patients receiving treatment		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of uncontrolled cases through NBS and treatment <sup>1</sup>	0	
Prevalence of uncontrolled RHD deficiency after newborn screening and treatment, per 1000 LB <sup>2</sup>	0	

LB = live births

NBS = newborn screening

RHD = Rhesus Haemolytic Disease of Newborn

\* If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

<sup>1</sup>Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

<sup>2</sup>Baseline birth prevalence – (Proportional reduction of uncontrolled cases of RHD X Baseline birth prevalence)



**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Needs Assessment Calculator 1: Quantitative baseline****Table RHD-NA1a Burden of Rhesus Haemolytic Disease of the Newborn in pregnancy, at birth and at population level**

Indicator	Chosen estimates			Notes
	Number (n)	n/1000 TB	Range of prevalence (/1000 TB)	
Annual affected live births (LB)	0	0	0	Drawn from sheet E2.4
Annual affected stillbirths (SB)	0	0	0	Drawn from sheet E2.4
Annual affected births (LB+SB)	0	0		Drawn from sheet E2.4
Annual affected persons (all age groups)	0	0	0	Drawn from sheet E1.1

**Table RCD-NA1b Rhesus Haemolytic Disease of the Newborn mortality indicators**

Indicator	Chosen estimates			Notes
	Number (n)	n/1000 LB	Range of prevalence (/1000 TB)	
Annual overall mortality	0			Drawn from sheet E3.4
Annual neonatal mortality	0	0	0	Drawn from sheet E3.4
Annual infant mortality	0	0	0	Drawn from sheet E3.4
Annual under-5 mortality	0	0	0	Drawn from sheet E3.4
Mean life expectancy at birth among affected people	0		0	Drawn from sheet E3.4

TB = total births (live births + stillbirths)

**Nigeria****Rhesus Haemolytic Disease of the Newborn****RHD Needs Assessment Calculator 3: Quantitative assessment of interventions**

<b>Table SCD-NA3a</b>	<b>Estimated prevalence in the absence of interventions for Sickle Cell Disease</b>	
Indicator	Number (n)	Prevalence (n/1000)
Potential live births		
Potential still births		

<b>Table RHD-NA3b</b>	<b>Current situation in relation to interventions before birth</b>		
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 LB
Effect of family planning, education			
Effect of anti D prophylaxis			
Effect of prenatal diagnosis and treatment			
Effect of neonatal diagnosis			
Effect of treatment of newborn			
Overall effect			

<b>Table RHD-NA3c</b>	<b>Target situation in relation to interventions before birth</b>		
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 LB
Effect of family planning, education			
Effect of anti D prophylaxis			
Effect of prenatal diagnosis and treatment			
Effect of neonatal diagnosis			
Effect of treatment of newborn			
Overall effect			

<b>Table RHD-NA3d</b>	<b>Current situation in relation to interventions after birth</b>		
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 LB
Newborn diagnosis			
Phototherapy			
Blood transfusion			
Social care and support			
Overall effect			

<b>Table RHD-NA3e</b>	<b>Target situation in relation to interventions after birth</b>		
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 LB
Newborn diagnosis			
Phototherapy			
Blood transfusion			
Social care and support			
Overall effect			

<b>Table RHD-NA3f</b>	<b>Current and desired outcomes</b>			
	<b>Current situation</b>		<b>Target situation</b>	
Indicator	Annual number (n)	Incidence (n/1000)	Annual number (n)	Incidence (n/1000)
<b>Estimated affected pregnancies</b>				
Live births (LB)	0	0		
Still births (SB)	0	0		
All births (LB+SB)	0	0		
<b>Estimated population prevalence</b>				
All age groups				
<b>Estimated mortality</b>				
Neonatal deaths	0	0		
Infant deaths	0	0		
Under-5 deaths	0	0		