## PHG Needs Assessment Calculator Thailand Congenital Heart Disease

Welcome to the PHG Health Needs Assessment Calculator for Congenital Heart Disease. The contents of this file are listed below.

Full name of the sheet	Short name
Country demographic, maternal health and socioeconomic indicators	Demography
Country health service data	HealthServices
CHD Epidemiology 1.1: Country epidemiology	CHD-E1.1
CHD Epidemiology 1.2: International comparison	CHD-E1.2
CHD Epidemiology 2.1: Data on affected pregnancies: Research studies	CHD-E2.1
CHD Epidemiology 2.2: Data on affected pregnancies: Surveillance	CHD-E2.2
CHD Epidemiology 2.3: Data on affected pregnancies: Other sources	CHD-E2.3
CHD Epidemiology 2.4: Summary of affected pregnancies	CHD-E2.4
CHD Epidemiology 2.5: Sub-population variation in affected pregnancies	CHD-E2.5
CHD Epidemiology 3.1: Mortality data: Research studies	CHD-E3.1
CHD Epidemiology 3.2: Mortality data: Vital registration data	CHD-E3.2
CHD Epidemiology 3.3: Mortality data: Other sources	CHD-E3.3
CHD Epidemiology 3.4: Summary mortality estimates	CHD-E3.4
CHD Epidemiology 3.5: Sub-population variation in mortality	CHD-E3.5
CHD Epidemiology 4.1: Population prevalence: Research studies	CHD-E4.1
CHD Epidemiology 4.2: Population prevalence: Other sources	CHD-E4.2
CHD Epidemiology 4.3: Summary of population prevalence	CHD-E4.3
CHD Epidemiology 4.4: Sub-population prevalence variation	CHD-E4.4
CHD Interventions 1: Effect of folic acid fortification	CHD-Interv1
CHD Interventions 2: Effect of folic acid supplementation	CHD-Interv2
CHD Interventions 3: Effects of prenatal screening and pregnancy management	CHD-Interv3
CHD Needs Assessment Calculator 1: Quantitative baseline	CHD-NA1
CHD Needs Assessment Calculator 3: Quantitative assessment of interventions	CHD-NA3

(There is no sheet CHD-NA2.)

Note: The Calculator sheets already contain modelled estimates from the PHGDB; note that these estimates do not include CHD associated with chromosomal disorders (e.g. Down's syndrome) and other non-cardiac malformations.

## Thailand Shared Data Demographic, maternal health and socio-economic indicators

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Demography information from that Calculator into here. The information should be the same.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Number of persons by age-group and sex		Estimates		Yo	our estimat	es	Cho	sen estima	ates
Age group	Male	Female	Total	Male	Female	Total	Male	Female	Total
0-4 years	1844815	1771822.8558	3616638			0			0
5-9 years	1544913	1480462.3473	3025375			0			0
10-14 years	1543617	1443636.0852	2987253			0			0
15-19 years	1476493	1357091.1013	2833584			0			0
20-24 years	1229710	1148159.0792	2377869			0			0
25-29 years	907707	887978.56398	1795686			0			0
30-34 years	692827.3	695264.65023	1388092			0			0
35-39 years	483223.3	501527.11796	984750.4			0			0
40-44 years	387883	411378.46007	799261.5			0			0
45-49 years	335165.6	358327.48264	693493.1			0			0
50-54 years	267219.2	278769.79026	545989			0			0
55-59 years	203903.8	196545.65274	400449.5			0			0
60-64 years	161471.4	151837.9478	313309.4			0			0
65+ years	376015.6	354270.43973	730286			0			0
Total	11454963	11037071.574	22492035	0	0	0	0	0	0
Female population aged 15-44 years		5001398.9728			-			-	
Data year		2009 report	ed in 2011						
Source, Year			UN 2011						

#### Ethnicity. Please enter data for the main ethnic groups if you are working with a population that is different from that of the country.

Ethnic group	Number	% population

Fertility and mortality	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Crude birth rate: live births (LB) / year / 1000 population	11.86	Unicef, 2013				loui
		,				
Still birth rate (SB): Still births (SB) / year / 1000 total	4.35	WHO, 2009				
births						
Total births in 1000s (LB+SB) per year	824	Unicef, 2013				
Infant mortality rate: infant deaths / 1000 LB / year	10.6	Unicef, 2013				
Under-5 mortality rate: U5 deaths / 1000 LB / year	12.3	Unicef, 2013				
Percentage births in women >35 years						
Life expectancy at birth (yrs)	74.13	Unicef, 2013				
% of marriages consanguineous						

	Estimate	Source, Year	Your	Source,	Chosen	Source,
Maternal health			estimate	Year	estimate	Year
Prenatal visits – at least 1 visit (%)	99.1	Unicef, 2013				
Prenatal visits – at least 4 visits (%)	79.6	Unicef, 2013				
Births attended by skilled health personnel (%)	99.5	Unicef, 2013				
Contraception prevalence rate (%)	79.6	Unicef, 2013				
Unmet need for family planning (%)	3.1	WHO, 2006				
Total fertility rate	1.56	Unicef, 2013				
% home births						
% births at health care services	99.40	Unicef, 2013				
	Estimate	Source, Year	Your	Source,	Chosen	Source,
Newborn health			estimate	Year	estimate	Year
Number of neonatal examinations by SBA / trained staff						
% neonatal examinations by SBA/ trained staff						

						Source,
Socio-economic indicators	Estimate	Source, Year	estimate	Year	estimate	Year
Gross national income per capita (PPP int. \$)	8390	Unicef, 2013				
% population living on < US\$1 per day	<2.0	Unicef, 2013				
Birth registration coverage (%)	99.4	WHO 2005-				
Death registration coverage (%)	50-74	WHO,2006				

LB = live births PPP = purchasing power parity SBA = skilled birth attendant PHG FOUNDATION

HealthServices

Thailand Shared Data Health Services Data

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Health Services information from that Calculator into here. The information should be the same.

This section provides health-service-related information for your country.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Health Expenditure	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Per capita total expenditure on health (PPP int. \$)	353.3	WHO 2011				
Total expenditure on health as percentage of GDP	4.1	WHO 2011				
Per capita government expenditure on health (PPP int. \$)	266.6	WHO 2011				
External resources for health as percentage of total expenditure on health	27	WHO 2011				
General government expenditure on health as percentage of total expenditure on health	75.5	WHO 2011				
Out-of-pocket expenditure as percentage of private expenditure on health	55.8	WHO 2011				
Private expenditure on health as percentage of total expenditure on health	24.5	WHO 2011				
General government expenditure on health as percentage of total government expenditure	14.5	WHO 2011				

		Source,	Your	Source,	Chosen	Source,
Health Workforce	Estimate	Year	estimate	Year	estimate	Year
Number of nursing and midwifery personnel	96704	WHO, 2004				
Nursing and midwifery personnel density (per 10,000 population)	15.2	WHO, 2004				
Number of physicians	18918	WHO, 2004				
Physician density (per 10,000 population)	2.98	WHO, 2004				
Number of obstetricians						
Number of paediatricians						
Number of paediatric surgeons						
Number of paediatric cardiac surgeons						
Number of paediatric neurosurgeons						
Number of clinical geneticists						
Number of genetic counsellors						
Number of community health workers						
Number of skilled birth attendants (SBA)						
Density of SBA						
Number of lab staff providing cytogenetic testing						
Number of lab staff providing molecular genetics						

Number of lab staff providing biochemical tests for genetics			
Number of skilled health attendants			

Infrastructure	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Number of maternity units						
Number of services providing specialised care for people with CD						
Number of family planning services						
Number of preconception services						
Number of services providing prenatal care						
Number of services providing newborn care						
Number of facilities providing genetic services						
Number of laboratories providing cytogenetics						
Number of laboratories providing molecular genetics						
Number of laboratories providing biochemical tests for genetics						
Number of facillities for safe terminations of pregnancies for fetal defects						

PPP = purchasing power parity GDP = gross domestic product

SBA = skilled birth attendant

CD = congenital disorders

## Thailand Congenital Heart Disease CHD Epidemiology 1.1: Country epidemiology

Epidemiological indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Year of estimate						
Prevalence at birth and by age-group (/1	000)					
Live birth prevalence (LB)			3.33			
Stillbirth prevalence (SB)			0.05			
Total birth prevalence (LB+SB)			3.38			
All age groups						
<1 year old						
1-4 year olds						
5-14 year olds						
15-44 year olds						
45+ year olds						
Number of cases by age group					·	
Annual live births			2,906			
All age groups						
<1 year olds						
1-4 year olds						
5-14 year olds						
15-44 year olds						
45+ year olds						
No. of cases by level of impairment					·	
No or minor disability						
Moderate disability*						
Severe disability*						
Mortality and morbidity					·	
Mean life expectancy (yrs)			48.5			
No. deaths < 1yr			607			
No. deaths 1-4 yrs			47			
No. deaths < 5 yrs			654			
Infant mortality / 1000 LB			0.70			
Under-5 mortality / 1000 LB			0.75			
Years of life lost						
LD live birther OD stillbirthe * Mademate						

LB = live births; SB = stillbirths \* Moderate = compensated cardiac problems, premature death preceded by average 1 year, and/or exertional disability. Severe = lifelong exertional disability

### Thailand

## Congenital Heart Disease

CHD: Epidemiology 1.2: International comparison

	Your chosen			
Epidemiological indicator	estimates	Country	Region	World
Prevalence at birth and by age-group (/1000 p	eople)		(Asia, Southeast)	
Live birth prevalence (LB)		3.33	3.17	3.24
Stillbirth prevalence (SB)		0.05	0.00	0.02
Total birth prevalence (LB+SB)		3.38	3.17	3.26
All age groups				
<1 year olds				
1-4 year olds				
5-14 year olds				
15-44 year olds				
45+ year olds				
Number of cases by age-group				
Annual live births		2,906	36,281	434,904
All age groups				
<1 year olds				
1-4 year olds				
5-14 year olds				
15-44 year olds				
45+ year olds				
No. cases by level of impairment				
No or minor disability				
Moderate disability*				
Severe disability*				
Mortality and morbidity				
Mean life expectancy (yrs)		48.5	33	29
No. deaths < 1yr		607	16,150	210,093
No. deaths 1-4 yrs		47	458	5,002
No. deaths < 5 yrs		654	16,608	215,094
Infant mortality / 1000 LB		0.70	0.45	0.48
Under-5 mortality / 1000 LB		0.75	0.46	0.49
Years of life lost				

LB = live births; SB = stillbirths \* Moderate = compensated cardiac problems, premature death preceded by average 1 year, and/or exertional disability. Severe = lifelong exertional disability

#### Thailand

#### Congenital Heart Disease

### CHD Epidemiology 2.1: Data on affected pregnancies: Research studies

Study author, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies listed above (or in section CHD-E2.1 of the Tool), enter the best estimates for the prevalence of affected births, still births and terminations in the country, and a range of values to reflect uncertainty or within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	LB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			
Estimates for the total country/territory	Number of affected stillbirths	SB prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			
Estimates for the total country/territory	Number of terminations of pregnancy due to condition	ТоР / 1000 ТВ	Comments
Best estimate			
Lower estimate			
Higher estimate			

## Thailand Congenital Heart Disease CHD Epidemiology 2.2: Data on affected pregnancies: Surveillance

Based on surveillance data, enter the best estimates for the prevalence of the condition in live births, stillbirths and pregnancy terminations in the country. Give a range of values to reflect uncertainty and within-country variation, and use comments for information on data quality, uncertainty and representativeness.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the total country/territory	Number of affected live births	Birth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

	Number of affected stillbirths	Stillbirth prevalence / 1000 TB	Comments
Best estimate			
Lower estimate			
Higher estimate			

Estimates for the total country/territory	Number of ToP due to condition	ТоР / 1000 ТВ	Comments
Best estimate			
Lower estimate			
Higher estimate			

# Thailand Congenital Heart Disease CHD Epidemiology 2.3: Data on affected pregnancies: Other sources

	Source 1:	Source 2:	Notes
Enter year and source of data – use last year with information available.			
Basic Numbers			
Number of affected live births / year, from data source			
Total number of live births / year, from data source			
Number of affected still births / year, from data source			
Total number of stillbirths / year, from data source			
Number of ToP for affected fetus / year from data source			
Total number of affected births / year (live and still)	(	0 0	
Total number of births / year, from data source	(	0 0	
Total number of ToP / year, from data source			
Total number of women aged 15-44			
Live birth prevalence: recorded and estimated			
Recorded live birth prevalence (affected recorded live births / 1000 recorded total births)	#DIV/0	! #DIV/0!	
Estimated completeness of recording: what proportion of true affected live births in your data source were recorded?	)		Range: 0 to 1
Estimated coverage of recorded live births (number of recorded live births / total live births in country or territory)			Range: 0 to 1
Estimated live birth prevalence (recorded prevalence / completeness)	#DIV/0	! #DIV/0!	
Estimated true number of affected live births in data source (number of recorded affected live births / completeness)	#DIV/0	! #DIV/0!	
Estimated number of affected live births in total population (number of affected live births from data source / (coverage x completeness))	#DIV/0	! #DIV/0!	
Stillbirth prevalence: recorded and estimated			
Recorded stillbirth prevalence (affected recorded still births / 1000 recorded total births)	#DIV/0	! #DIV/0!	
Estimated completeness of recording: what proportion of true affected stillbirths in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded stillbirths (number of recorded still births / total still births in country or territory)			Range: 0 to 1
Estimated stillbirth prevalence (recorded prevalence / completeness)	#DIV/0	! #DIV/0!	
Estimated true number of affected stillbirths in data source (number of recorded affected still births / completeness)	#DIV/0	! #DIV/0!	
Estimated number of affected stillbirths in total population (number of affected still births from data source / (coverage x completeness))	#DIV/0	! #DIV/0!	

ToP prevalence: recorded and estimated			
Recorded ToP prevalence (ToP in affected fetuses / 1000 women aged 15-44)	#DIV/0!	#DIV/0!	
Estimated completeness of recording: what proportion of true affected pregnancy terminations in your data source were recorded?			Range: 0 to 1
Estimated coverage of recorded ToP (number of recorded ToP / total ToP in country or territory)			Range: 0 to 1
Estimated ToP prevalence (recorded prevalence / estimated completeness)	#DIV/0!	#DIV/0!	
Estimated true number of ToP in data source (number of recorded ToP / completeness)	#DIV/0!	#DIV/0!	
Estimated number of ToP in total population (number of ToP from data source / (coverage x completeness))	#DIV/0!	#DIV/0!	

**Based on the sources above**, enter the best prevalence estimates for your population, and a range of values to reflect uncertainty of estimates and within country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Estimates for the whole country/territory	Number of affected live births	LB prevalence / 1000 TB
Best estimate		
Lower estimate		
Higher estimate		
Estimates for the whole country/territory	Number of affected still births	SB prevalence / 1000 TB
Best estimate		
Lower estimate		
Higher estimate		
Estimates for the whole country/territory	Number of ToP due to condition	ТоР /1000 ТВ
Best estimate		
Lower estimate		
Higher estimate		

## Thailand Congenital Heart Disease CHD Epidemiology 2.4: Summary of affected pregnancies

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Number of annual affected live births			2,906			
Annual birth prevalence / 1000 TB			3.33			
Number of annual affected still births			3			
Stillbirth prevalence / 1000 TB/year			0.05			
Number of <b>terminations of</b> <b>pregnancy</b> in affected fetuses /year						
Affected ToP / 1000 TB						

If there are specific sub-types of condition, you can repeat this exercise below. However, you should consider (a) whether subtypes would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

# Thailand Congenital Heart Disease CHD Epidemiology 2.5: Sub-population variation in affected pregnancies

If the birth prevalence rates vary by population sub-group (e.g. geographically or by another factor), indicate any population groups with different prevalence estimates from the whole population and describe reasons for variation. If a group is substantially different from the general population, you may wish to conduct a needs assessment for that group alone.

Population sub- group	Number of affected live births	LB prevalence / 1000 TB	Reason for variation

Population sub- group	Number of affected stillbirths	SB prevalence / 1000 TB	Reason for variation

•	Number of ToP in affected pregnancies	ToP prevalence / 1000 TB	Reason for variation

## Thailand Congenital Heart Disease CHD Epidemiology 3.1: Mortality data: Research studies

Source, year, site	Sample size	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for the specific mortality by age-group e.g. infant, under-5s, etc., as appropriate, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

Mortality estimates	Number of deaths	Ratio (deaths / 1000 LB)	Comments
Neonatal group (<28 days)			
Best estimate			
Lower estimate			
Higher estimate			
Infant group (<1 year)			
Best estimate			
Lower estimate			
Higher estimate			
Under-5 group (<5 years)			
Best estimate			
Lower estimate			
Higher estimate			
Other age group:			
Best estimate			
Lower estimate			
Higher estimate			

LB = live births

## Thailand

### Congenital Heart Disease

CHD Epidemiology 3.2: Mortality data: Vital registration data

Fill in the blank cells based on your vital registration data.		
Enter year and source of data		
Registered data		
Total registered live births		
Registered condition-specific neonatal deaths (first 28 days of life)		
Registered condition-specific infant deaths (first year of life)		
Registered condition-specific under-5 deaths (first 5 years of life)		
Registered condition-specific neonatal mortality ratio (condition-specific neonatal deaths / 1000 live births in the same year)	#DIV/0!	
Registered condition-specific infant mortality (condition-specific infant deaths / 1000 live births in the same year)	#DIV/0!	
Registered condition-specific under-5 mortality (condition-specific under-5 deaths / 1000 live births in the same year)	#DIV/0!	

Adjustment for under-ascertainment of cause of death and sub-registration of deaths: Enter estimates in the highlighted cells. It is not always possible to adjust the estimates, in which case you may give the value '1', accepting that the estimates in these cases will usually be biased towards low values. (Or you may move to the next section.) It is assumed that under-ascertainment is stable across age-groups; if ascertainment varies by age-group, you could use separate estimates for each age group.

Estimated completeness of recording: what proportion of deaths in affected persons were registered as such?		Range: 0 to 1
Population coverage: what proportion of the total country/territory population is covered by the vital registration?		Range: 0 to 1
Death ascertainment (population coverage x completeness)	0	
Estimated values for the total country/ territory population		
Estimated number of live births in total population	#DIV/0!	
Estimated number of neonatal deaths in total population	#DIV/0!	
(number of deaths registered in neonatal period / ascertainment)		
Estimated number of infant deaths in total population	#DIV/0!	
(number of deaths registered in first year of life / ascertainment)		
Estimated number of under-5 deaths in total population	#DIV/0!	
(number of deaths registered in under-5s / ascertainment)		
Estimated neonatal mortality ratio (estimated neonatal deaths / 1000 live births)	#DIV/0!	
Estimated infant mortality ratio (estimated infant deaths / 1000 live births)	#DIV/0!	
Estimated under-5 mortality ratio (estimated under-5 deaths / 1000 live births)	#DIV/0!	

### Thailand Congenital Heart Disease CHD Epidemiology 3.3: Mortality data: Other sources

Source, year, site	Sample size	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population. If studies are not representative of the national population you may need to weight your data (see the Guide

for explanation on weighting and help with the calculations).

	Neonatal mortality		Infant mortality		Under-5 mortality	
Estimates for the total country/territory	Value	Ratio/1000 LB	Value	Ratio/1000 LB	Value	Ratio/1000 LB
Best estimate						
Lower estimate						
Higher estimate						

### Thailand Congenital Heart Disease CHD Epidemiology 3.4: Summary mortality estimates

Indicator	Your estimates	Range	PHGDB minimum estimates	Chosen estimates	Range	Source
Year of data collection						
Number of annual deaths in affected persons						
Number of annual live births (in 1000s)			873			
Number of annual affected neonatal deaths			379			
Number of affected neonatal deaths / 1000 LB			0.43			
Number of annual affected infant deaths			607			
Number of affected infant deaths / 1000 LB			0.70			
Number of annual affected under-5 deaths	1		654			
Number of affected under-5 deaths / 1000 LB			0.75			
Mean life expectancy at birth in affected people			48.5			
Other indicators (e.g. survival following surgical procedure, etc)						

#### Thailand

# Congenital Heart Disease

CHD Epidemiology 3.5: Sub-population variation in mortality

Age group: neonatal Population sub-group	Cause-specific, group-specific neonatal mortality ratio / 1000 LB	Reason for variation

Age group: infant Population sub-group	Cause-specific, group-specific infant mortality ratio / 1000 LB	Reason for variation

		Cause-specific, group-specific	Reason for variation
Population sub-group	affected persons	under-5 mortality ratio / 1000 LB	

Age group:		Cause-specific, group-specific	Reason for variation	
Population sub-group	affected persons	mortality ratio / 1000 population		

# Thailand Congenital Heart Disease CHD Epidemiology 4.1: Population prevalence: Research studies

Study, year, site	Study quality and representativeness	Main findings

Based on the studies above, enter the best estimates for population prevalence, and a range of values to reflect uncertainty of estimates and within-country variation.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

## Thailand Congenital Heart Disease CHD Epidemiology 4.2: Population prevalence: Other sources

Source, year, site	Data quality and representativeness	Main findings

Based on data from the sources above, enter estimates for the disease-specific deaths and mortality rates in your population.

If studies are not representative of the national population you may need to weight your data (see the Guide for explanation on weighting and help with the calculations).

	Prevalence / 1000 persons	Range	Comments
Best estimate			
Lower estimate			
Higher estimate			

## Thailand Congenital Heart Disease CHD Epidemiology 4.3: Summary of population prevalence

Source of estimates	Estimated total population number of affected persons	Range	Estimated total population prevalence / 1000 persons	Range
1				
2				
3				
4				
5				
PHGDB				
Chosen estimates				

## Thailand Congenital Heart Disease CHD Epidemiology 4.4: Sub-population prevalence variation

Population sub-group	Number of affected people	Population prevalence per 1000 people	Reason for variation
		#DIV/0!	

If there are specific sub-types of condition, you can repeat this exercise (copy table and paste below). However, you should consider (a) whether sub-types would have different implications for advocacy, and (b) whether a sub-type might require a full, specific needs assessment.

Formula in column D: Number of affected people/ (Total number of people in population subgroup/1000)

## Thailand Congenital Heart Disease CHD Interventions 1: Effect of folic acid fortification

This sheet allows you to estimate the potential reduction in CHD prevalence through fortification of food with folic acid. Please start by entering values reflecting your current situation. If you have no fortification programme, enter 0 for coverage. Below, you may adjust dosage and coverage levels to demonstrate the effects of different intervention scenarios.

Current situation	Notes
Present estimated CHD prevalence per 1000 TB	
Present dosage (ppm)	Range: 1.5 to 3
Present coverage of fortification	Range: 0 to 1
Baseline CHD prevalence per 1000 TB, with no folic acid fortification*1	

Potential scenarios, based on your present situation		
Vary dosage (ppm)	Range: 1.5 to 3	
Vary proportional population coverage Estimated reduction in CHDs through folic acid fortification, per 1000	Range: 0 to 1	
Estimated reduction in CHDs through folic acid fortification, per 1000 TB <sup>2</sup>	0.000 Do not delete this value!	
Resulting prevalence of CHDs after folic acid fortification, per 1000 TB <sup>3</sup>	0.000 Do not delete this value!	

ppm = parts per million

TB = total births (live births + still births)

\* The effect of folic acid on CHD is assumed to be 25% of the effect on neural tube defects.

The regression formula underlying the effect on neural tube defects is given in the NTD Calculator in this Toolkit.

\*\* Not considering the effects of other interventions on prevalence.

<sup>1</sup>(Present estimated prevalence-(1.07\*coverage\*0.25)+(0.15\*ppm\*coverage\*0.25))/(1-0.88\*coverage\*0.25))) <sup>2</sup>((0.25\*(Baseline CHD-(1.07\*coverage+0.12\*baseline CHD\*coverage-0.15\*dosage\*coverage+baseline-baseline\*coverage))))

<sup>3</sup>Baseline CHD prevalence – estimated reduction in CHD after fortification

### Thailand Congenital Heart Disease CHD Interventions 2: Effect of folic acid supplementation

This sheet allows you to estimate the potential reduction in CHD incidence through folic acid supplementation for pregnant women. Please enter a value for population coverage of folic acid supplementation, to determine its potential effect.

Effect of supplementation (with no fortification)		Notes	
Baseline prevalence with no folic acid intervention (per 1000 TB)		This can be taken from the appropriate cell in the previous sheet.	
Maximum proportional reduction (assuming 100%			
coverage)	0.18	This value is fixed at 0.18	
Population supplementation coverage		Range: 0 to 1	
Actual proportional reduction	0	Maximum proportional reduction x Coverage	
Actual prevalence reduction (per 1000 TB)	0.000	Baseline prevalence x Actual proportional reduction	

New prevalence		Baseline prevalence -((Maximum prop. Reduction x Population supplementation coverage) x Baseline prevalence))
% prevalence reduction	#DIV/0!	1-(New prevalence/Baseline prevalence)
Absolute prevalence reduction (per 1000 TB)	0.000	Baseline prevalence -New prevalence

Now you can see below the potential combined effect of folate fortification and supplementation:

Additional effect of supplementation, given fortification	0.1 This value can be changed.

	New prevalence	
After fortification		This can be taken from the appropriate cell above
After supplementation	0.000	Same as new prevalence
After fortification and supplementation		Prevalence after fortification-(Additional effect of supplementation*prevalence after supplementation)

### TB = total births (live births + still births)

CHD = congenital heart disease

<sup>1</sup>New Prevalence after fortification-(Additional effect of supplementation x Final prev. following supplemen.)

<sup>2</sup>If New prevalence after fortification < minimum prevalence then use (Baseline prev – min prevalne)/baseline prevalance)

Otherwise use: (Baseline prevalence – new prevalence after fortification and supplementation)/baseline prevalence

# Thailand Congenital Heart Disease CHD Interventions 3: Effect of prenatal screening and pregnancy management

Baseline prevalence: fetuses affected by CHD, per 1000 TB Variables		See previous sheet. Use baseline prevalence either before or after folic acid fortification and supplementation.
Coverage of prenatal screening		Range: 0 to 1
Proportion of diagnosed pregnancies receiving treatment*		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of uncontrolled cases of CHD through PNS and treatment <sup>1</sup>	0%	
Prevalence of uncontrolled CHD after PNS and treatment, per 1000 total births <sup>2</sup> 0.000		
Final prevalence: affected live births after PNS & treatment, per 1000 total births <sup>3</sup>		

PNS = prenatal screening

TB = total births (live births + still births)

CHD = congenital heart disease

\*Treatment in this case refers to diagnosis and appropriate management of pregnancy

<sup>1</sup>Coverage of screening X Proportion of diagnosed pregnancies receiving treatment X Effectiveness of treatment <sup>2</sup>Proportional reduction of uncontrolled cases x Baseline prevalence <sup>3</sup>Baseline prevalence – prevalence of uncontrolled CHD

#### CHD Interventions: Effect of prenatal screening and pregnancy termination

Assumption: prenatal services are equally used for cases which would lead to still births and live births. This could overestimate the impact of ToP if in fact ToP is more likely for severe cases that would result in stillbirth. Conversely, the impact of ToP could be underestimated if screening is only available to high-income women at lower risk. 100% specificity of prenatal diagnosis assumed.

Baseline prevalence, per 1000 TB (LB + SB)		See previous two sheets. Use baseline either before or after folic acid interventions.
Variables		
Coverage of prenatal screening		Range: 0 to 1
Proportion of screen-positive cases receiving diagnosis		Range: 0 to 1
Proportion of diagnosed cases ending in pregnancy termination		Range: 0 to 1
Results		
% prevalence reduction due to PND & pregnancy termination <sup>1</sup>	0%	
Prevalence reduction due to PND & pregnancy termination, per 1000 TB <sup>2</sup>	0.000	
Final birth prevalence of CHDs after PND & pregnancy termination, per 1000 TB <sup>3</sup>	0.000	

PNS = prenatal screening

ToP = termination of pregnancy

TB = total births (live births + still births)

<sup>1</sup>Coverage of screening X Proportion of screen-positive cases receiving diagnosis x Proportion of cases ending in pregnancy termination <sup>2</sup>% prevalence reduction due to PND and termination x Baseline prevalence <sup>3</sup>Baseline prevalence – Prevalence reduction due to PND & termination

# Thailand Congenital Heart Disease CHD Needs Assessment Calculator 1: Quantitative baseline

# Table CHD-NA1a Burden of Neural Tube Defects in pregnancy, at birth and at population level

		Chosen estimates		
Indicator	Number (n)		Range of prevalence (/1000 TB)	
Annual affected live births (LB)	(	0 0	C	Drawn from sheet E2.4
Annual affected stillbirths (SB)	(	0 0	C	Drawn from sheet E2.4
Annual affected births (LB+SB)	(	0 0		Drawn from sheet E2.4
Annual affected persons (all age	(	0 0	C	Drawn from sheet E1.1
groups)		·	•	·

# Table CHD-NA1b Neural Tube Defects mortality indicators

	Chosen estimates			Notes
Indicator	Number (n)	1	Range of prevalence (/1000 TB)	
Annual overall mortality	C			Drawn from sheet E3.4
Annual neonatal mortality	C	0	0	Drawn from sheet E3.4
Annual infant mortality	C	0	0	Drawn from sheet E3.4
Annual under-5 mortality	C	0	0	Drawn from sheet E3.4
Mean life expectancy at birth among affected people	C		0	Drawn from sheet E3.4

TB = total births (live births + stillbirths)

#### Thailand

### Congenital Heart Disease

### CHD Needs Assessment Calculator 3: Quantitative assessment of interventions

	Estimated prevalence in the absence of interventions for Neural Tube Defects		
Indicator	Number (n)	Prevalence (n/1000)	
Potential live births			
Potential still births			

Table CHD-NA3b	Current situation in relation to interventions before birth			
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 TB	
Effect of family planning, education				
Effect of multivitamin and folic acid supplementation				
Control of teratogens				
Management of diseases in pregnancy, e.g diabetes				
Rubella prevention				
Effect of prenatal diagnosis				
Overall effect				

Table CHD-NA3c	Target situation in relation	on to interventions before	birth
Intervention	Coverage (%)	Cases averted (n)	Cases averted/1000 TB
Effect of family planning, education			
Effect of multivitamin and folic acid supplementation			
Control of teratogens			
Management of diseases in pregnancy, e.g diabetes			
Rubella prevention			
Effect of prenatal diagnosis			
Overall effect			

Table CHD-NA3d	Current situation in rela	ation to interventions afte	r birth
Intervention	Coverage (%)	Cases managed (n)	Cases managed/1000 TB
Effect of newborn screening			
Effect of newborn diagnosis			
Effect of echocardiography services			
Effect of primary care diagnosis			
Effect of paediatric cardiology			
Effect of paediatric cardiac surgery			
Effect of acute clinical care			
Effect of social care and support			
Effect of rehabilitation			
Overall effect			

Target situation in relation	on to interventions after	birth	
Coverage (%)	Cases managed (n)	Cases managed/1000 TB	1
			_
			-
			_
			-
			-
			1
			1
			1
			1
Current and desired out	comes		
Current situation		Target situation	
Annual number (n)	Prevalence (n/1000)	Annual number (n)	Prevalence (n/1000)
0		0	
0		0	
			i
0		0	
C	0	0	
C		0	
0		0	
		0	
15			
	Coverage (%) Current and desired out Current situation Annual number (n) Current Current	Coverage (%)       Cases managed (n)         Cases managed (n)       Image: Cases managed (n)         Image: Cases managed (n)       Imaged (n)     <	Current and desired outcomes       Current situation       Annual number (n)       Prevalence (n/1000)       Annual number (n)

TB = total births (live births + stillbirths)