PHG Needs Assessment Calculator Congo Newborn screening

Welcome to the PHG Health Needs Assessment Calculator for Newborn Screening. The contents of this file are listed below.

Full name of the sheet	Short name
Country demographic, maternal health and socioeconomic indicators	Demography
Country health-service indicators	HealthServices
Existing screening programmes for congenital disorders	NBS-NA1.1
Details of newborn screening programmes	NBS-NA1.2
Effect of newborn screening and treatment on congenital hypothyroidism	NBS-CHT
Effect of newborn screening and treatment on G6PD deficiency	NBS-G6PD
Effect of newborn screening and treatment on Rhesus haemolytic disease of newborn	NBS-RHD
Effect of newborn screening and management on sickle cell disease	NBS-SCD
Effect of newborn screening and management on thalassaemias	NBS-THAL
Effect of newborn screening and treatment on orofacial clefts	NBS-OFC
Effect of newborn screening and treatment on phenylketonuria	NBS-PKU
Effect of newborn screening and management on cystic fibrosis	NBS-CF

Congo

Shared Data

Demographic, maternal health and socio-economic indicators

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Demography information from that Calculator into here. The information should be the same.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Number of persons by age-group and sex		Estimates		Your estimates		Cho	sen estima	ates	
Age group	Male	Female	Total	Male	Female	Total	Male	Female	Total
0-4 years	39737	29729	77147			0			0
5-9 years	33771	32005	65776			0			0
10-14 years	31576	29738	61314			0			0
15-19 years	39899	38014	77913			0			0
20-24 years	54299	51864	106163			0			0
25-29 years	51929	50775	102704			0			0
30-34 years	47095	46415	93510			0			0
35-39 years	45889	46709	92598			0			0
40-44 years	42306	45210	87516			0			0
45-49 years	43536	48552	92088			0			0
50-54 years	42675	50409	93084			0			0
55-59 years	37982	48391	86373			0			0
60-64 years	31689	43781	75470			0			0
65+ years	75100	153281	228381			0			0
Total	0	0	1340160	0	0	0	0	0	0
Female population aged 15-44 years		0			0			0	
Data year		2010 report	ed in 2011	·					
Source, Year			UN 2011						

Ethnicity. Please enter data for the main ethnic groups if you are working with a population that is different from that of the country.

Ethnic group	Number	% population

Fertility and mortality	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Struderthirtateate: dives hirtherths By by 9 and a 1900 bonulation	42.90	Unicef, 2013				
births	28.56	WHO, 2009				
Total births in 1000s (LB+SB) per year	2912	Unicef, 2013				
Infant mortality rate: infant deaths / 1000 LB / year	110.60	Unicef, 2013				
Under-5 mortality rate: U5 deaths / 1000 LB / year	167.70	Unicef, 2013				
Percentage births in women >35 years						
Life expectancy at birth (yrs)	48.40	Unicef, 2013				
% of marriages consanguineous						

Maternal health	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Prenatal visits – at least 1 visit (%)	88.8	Unicef, 2013				
Prenatal visits – at least 4 visits (%)	44.7	Unicef, 2013				
Births attended by skilled health personnel (%)	80.4	Unicef, 2013				
Contraception prevalence rate (%)	17.3	Unicef, 2013				
Unmet need for family planning (%)	16.2	WHO, 2005				
Total fertility rate	5.66	Unicef, 2013				
% home births						
% births at health care services	74.90	Unicef, 2013				
Newborn health	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Number of neonatal examinations by SBA / trained staff						
% neonatal examinations by SBA/ trained staff						

			Your	Source, Year	Chosen	Source,
Socio-economic indicators	Estimate	Source, Year	estimate		estimate	Year
Gross national income per capita (PPP int. \$)	350	Unicef, 2013				
% population living on < US\$1 per day	54.1	Unicef, 2013				
Birth registration coverage (%)	27.8	WHO 2010				
Death registration coverage (%)						

LB = live births

PPP = purchasing power parity

SBA = skilled birth attendant

Congo Shared Data Health Services Data

Please read first! If you have already completed a needs assessment for a different topic in this country, you will be able to copy the Health Services information from that Calculator into here. The information should be the same.

This section provides health-service-related information for your country.

By default, the Toolkit contains information at the national level.

If you would like to use a different population, then replace country information with that of your specific population of interest.

Health Expenditure	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Per capita total expenditure on health (PPP int. \$)	32.1	WHO 2011				
Total expenditure on health as percentage of GDP	8.5	WHO 2011				
Per capita government expenditure on health (PPP int. \$)	10.8	WHO 2011				
External resources for health as percentage of total expenditure on health	31.4	WHO 2011				
General government expenditure on health as percentage of total expenditure on health	33.7	WHO 2011				
Out-of-pocket expenditure as percentage of private expenditure on health	65.7	WHO 2011				
Private expenditure on health as percentage of total expenditure on health	66.3	WHO 2011				
General government expenditure on health as percentage of total government expenditure	10.8	WHO 2011				

		Source,	Your	Source,	Chosen	Source,
Health Workforce	Estimate	Year	estimate	Year	estimate	Year
Number of nursing and midwifery personnel	3492	WHO, 2007				
Nursing and midwifery personnel density (per 10,000 population)	8.2	WHO, 2007				
Number of physicians	401	WHO, 2007				
Physician density (per 10 000 population)	0.95	WHO, 2007				
Number of obstetricians						
Number of paediatricians						
Number of paediatric surgeons						
Number of paediatric cardiac surgeons						
Number of paediatric neurosurgeons						
Number of clinical geneticists						
Number of genetic counsellors						
Number of community health workers						
Number of skilled birth attendants (SBA)						
Density of SBA						

Number of lab staff providing cytogenetic testing			
Number of lab staff providing molecular genetics			
Number of lab staff providing biochemical tests for genetics			
Number of skilled health attendants			

Infrastructure	Estimate	Source, Year	Your estimate	Source, Year	Chosen estimate	Source, Year
Number of maternity units	Louinato	Tour	Commuto	loui	Commuto	Tour
Number of services providing specialised care for people with CD						
Number of family planning services						
Number of preconception services						
Number of services providing prenatal care						
Number of services providing newborn care						
Number of facilities providing genetic services						
Number of laboratories providing cytogenetics						
Number of laboratories providing molecular genetics						
Number of laboratories providing biochemical tests for genetics						
Number of facillities for safe terminations of pregnancies for fetal defects						

PPP = purchasing power parity GDP = gross domestic product SBA = skilled birth attendant

CD = congenital disorders

Congo Newborn screening Existing screening programmes for congenital disorders

Condition	Tick if NBS programme exists	Tick if included in physical examination	Indicate whether NBS is provided at national or subnational level	Condition prevalence per 1000 newborns	Prevalence variation and high-risk populations
Eye problems					
Signs of heart disease					
Developmental dysplasia of hips					
Genital anomalies (e.g. undescended testicles)					
Orofacial clefts					
Dysmorphologies					
Hearing loss					
Congenital hypothyroidism					
G6PD deficiency					
PKU					
Cystic fibrosis					
Thalassaemias					
Sickle cell disease					
MCADD					
CAH					
Other					

NBS = newborn screening
G6PD = glucose-6-phosphate dehydrogenase
PKU = phenylketonuria
CAH= congenital adrenal hyperplasia
MCADD = medium-chain acyl-CoA dehydrogenase deficiency

Congo Newborn screening Details of newborn screening programmes

Condition	Age at screen	Coverage (%)	Coverage variation and high-risk populations	Estimated proportion of affected newborns detected	Target coverage (%)
Newborn physical ex	amination	1			
Basic examination*					
Examination for gross abnormalities*					
Detailed physical examination					
Newborn hearing scr	eening				
Crude screening					
Equipment based screening					
Newborn bloodspot s	screening				
Congenital hypothyroidism					
PKU					
Cystic fibrosis					
Sickle cell disease					
G6PD deficiency					
MCADD					
CAH					
Other					

PKU = phenylketonuria

G6PD = glucose-6-phosphate dehydrogenase

MCADD = medium-chain acyl-CoA dehydrogenase deficiency

CAH= congenital adreanal hyperplasia

^{*} As defined in the Background document section titled Newborn Screening Tests

Congo Newborn screening Effects of NBS and treatment on congenital hypothyroidism

Baseline birth prevalence of CHT, per 1000 total births*		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of positive-screened patients receiving diagnosis treatment		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of uncontrolled cases of CHT through NBS and treatment ¹	0	
Prevalence of uncontrolled CHT after newborn screening and treatment, per 1000 total births²	0	

LB = live births

CHT = congenital hypothyroidism

NBS = newborn screening

^{*} If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of uncontrolled cases of CHT X Baseline birth prevalence)

Congo Newborn screening Effects of NBS and treatment on G6PD deficiency

Baseline birth prevalence of G6PD deficiency, per 1000 LB		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of positive-screened patients receiving treatment		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of uncontrolled cases through NBS and treatment ¹	0	
Prevalence of uncontrolled G6PD deficiency after newborn screening and treatment, per 1000 LB ²	0	

LB = live births

NBS = newborn screening

G6PD = glucose-6-phosphate dehydrogenase

^{*} If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of uncontrolled cases of G6PD X Baseline birth prevalence)

Congo Newborn screening Effects of NBS and treatment on RHD

Baseline birth prevalence of RHD, per 1000 LB		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of positive-screened patients receiving treatment		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of uncontrolled cases through NBS and treatment ¹	0	
Prevalence of uncontrolled RHD deficiency after newborn screening and		
treatment, per 1000 LB ²	0	

LB = live births

NBS = newborn screening

RHD = Rhesus Haemolytic Disease of Newborn

^{*} If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of uncontrolled cases of RHD X Baseline birth prevalence)

Congo Newborn screening

Effects of NBS and management on sickle cell disease

Baseline birth prevalence of sickle cell disease, per 1000 LB		
Variables		
Proposed on the series of the		Range: 0 to 1
management		Range: 0 to 1
Effectiveness of management		Range: 0 to 1
Results		
Proportional reduction in unmanaged cases of SCD through NBS and treatment ¹	0	
Prevalence of unmanaged sickle cell disease after newborn screening and treatment, per 1000 LB ²	0	

LB = live births

SCD = sickle cell disease

NBS = newborn screening

^{*} If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of unmanaged cases of SCD X Baseline birth prevalence)

Congo Newborn screening Effects of NBS and management on thalassaemias

Baseline birth prevalence of thalassaemias, per 1000 LB		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of screen-positive patients referred for treatment		Range: 0 to 1
Effectiveness of management		Range: 0 to 1
Results		
Proportional reduction of prevalence of unmanaged thalassaemias through NBS and treatment ¹	0	
Prevalence of unmanaged thalassaemias after newborn screening and treatment, per 1000 LB ²	0	

LB = live births

NBS = newborn screening

* If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of unmanaged cases of thalassaemia X Baseline birth prevalence)

Congo Newborn screening Effects of NBS and treatment on orofacial clefts

Baseline birth prevalence of orofacial clefts, per 1000 LB		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of screen-positive patients receiving treatment		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of prevalence of untreated OFCs through NBS and treatment ¹	0	
Prevalence of untreated OFCs after newborn screening and treatment, per 1000 LB ²	0	

LB = live births

OFCs = orofacial clefts

NBS = newborn screening

^{*} If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of untreated cases of OFC X Baseline birth prevalence)

Congo Newborn screening Effects of NBS and treatment on phenylketonuria

D 1: 1: 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		
Baseline birth prevalence of PKU, per 1000 LB		
Variables		
Coverage of newborn screening		Range: 0 to 1
Proportion of positive-screened patients receiving treatment		Range: 0 to 1
Effectiveness of treatment		Range: 0 to 1
Results		
Proportional reduction of prevalence of clinical cases of PKU through NBS and treatment ¹	0	
Prevalence of symptomatic PKU after newborn screening and treatment, per 1000 LB ²	0	

LB = live births

PKU = phenylketonuria

NBS = newborn screening

¹Coverage of newborn screening X Proportion of screen-positive cases receiving treatment X Effectiveness of treatment

²Baseline birth prevalence – (Proportional reduction of prevalence of clinical cases of PKU X Baseline birth prevalence)

^{*} If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

Congo Newborn screening Effects of NBS and management on cystic fibrosis

Baseline birth prevalence of cycstic fibrosis, per 1000 LB		
Variables		
คิกษาสมาคา of เกืองเมลา เลืองเลืองเลืองเลืองเลืองเลืองเลืองเลือง		Range: 0 to 1
management		Range: 0 to 1
Effectiveness of management		Range: 0 to 1
Results		
Proportional reduction of prevalence of unmanaged cystic fibrosis through NBS and treatment ¹	0	
Prevalence of unmanaged cystic fibrosis after newborn screening and treatment, per 1000 LB ²	0	

LB = live births

NBS = newborn screening

* If you don't have data on birth prevalence but do have data on screening, you can estimate birth prevalence by combining the proportion screened positive with the number of total births. (This assumes that screening is randomly distributed in the population).

¹Coverage of newborn screening X Proportion of positive-screened patients referred for management X Effectiveness of management

²Baseline birth prevalence – (Proportional reduction of prevalence of unmanaged cases X Baseline birth prevalence)